

CHOLINE BINDING PROTEINS FOR ANTI-PNEUMOCOCCAL VACCINES

The research leading to the present invention was supported, in part, by Grant Nos.
5 AI 36445 and AI 38446 from the National Institutes of Health. Accordingly, the
Government may have certain rights in the invention.

The present application claims priority to Provisional Patent Application Serial No.
60/016,632, filed May 1, 1996, pursuant to 35 USC 119(e), the disclosure of which
10 is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates generally to choline binding proteins, methods for
15 isolating choline binding proteins, and the genes encoding such proteins. The
invention also relates to acellular vaccines to provide protection from bacterial
infection using such proteins, and to antibodies against such proteins for use in
diagnosis and passive immune therapy. In particular, the choline binding proteins
of the invention are useful as vaccines against pneumococcus. Where a choline
20 binding protein demonstrates activity as an adhesion protein, it is also useful as a
competitive inhibitor of bacterial adhesion, or to discover small molecule
antagonists of adhesion.

BACKGROUND OF THE INVENTION

25 Antibacterial Vaccine Strategies

Exported proteins in bacteria participate in many diverse and essential cell functions
such as motility, signal transduction, macromolecular transport and assembly, and
the acquisition of essential nutrients. For pathogenic bacteria, many exported
proteins are virulence determinants that function as adhesions to colonize and thus
30 infect the host or as toxins to protect the bacteria against the host's immune system
[International Patent Publication No. WO 95/06732, published March 9, 1995 by
Masure et al., which is specifically incorporated herein by reference in its entirety,

for a review, see Hoepelman and Tuomanen, *Infect. Immun.*, **60**:1729-33 (1992)].
However, other exported proteins may not directly mediate adhesion.

Since the development of the smallpox vaccine by Jenner in the 18th century,
5 vaccination has been an important armament in the arsenal against infectious
microorganisms. Prior to the introduction of antibiotics, vaccination was the major
hope for protecting populations against viral or bacterial infection. With the advent
of antibiotics in the early 20th century, vaccination against bacterial infections
became much less important. However, the recent resurgence of antibiotic-resistant
10 strains of infectious bacteria has resulted in the reestablishment of the importance of
anti-bacterial vaccines.

One possibility for an anti-bacterial vaccine is the use of killed or attenuated
bacteria. However, there are several disadvantages of whole bacterial vaccines,
15 including the possibility of a reversion of killed or attenuated bacteria to virulence
due to incomplete killing or attenuation and the inclusion of toxic components as
contaminants.

Another vaccine alternative is to immunize with the bacterial carbohydrate capsule.
20 Presently, vaccines against *Streptococcus pneumoniae* employ conjugates composed
of the capsules of the 23 most common serotypes of this bacterium. these vaccines
are ineffective in individuals most susceptible to pathological infection -- the young,
the old, and the immune compromised -- because of its inability to elicit a T cell
immune response. A recent study has shown that this vaccine is only 50%
25 protective for these individuals [Shapiro et al., *N. Engl. J. Med.* **325**:1453-60
(1991)].

An alternative to whole bacterial vaccines are acellular vaccines or subunit vaccines
in which the antigen includes a bacterial surface protein. These vaccines could
30 potentially overcome the deficiencies of whole bacterial or capsule-based vaccines.

Moreover, given the importance of exported proteins to bacterial virulence, these proteins are an important target for therapeutic intervention. Of particular importance are proteins that represent a common antigen of all strains of a particular species of bacteria for use in a vaccine that would protect against all strains of the bacteria. However, to date only a small number of exported proteins of Gram positive bacteria have been identified, and none of these represent a common antigen for a particular species of bacteria.

Recently, apparent fusion proteins containing PhoA were exported in species of Gram positive and Gram negative bacteria (Pearce and Masure, 1992, *Abstr. Gen. Meet. Am. Soc. Microbiol.* 92:127, abstract D-188). This abstract reports insertion of pneumococcal DNA upstream from the *E. coli phoA* gene lacking its signal sequence and promoter in a shuttle vector capable of expression in both *E. coli* and *S. pneumoniae*, and suggests that similar pathways for the translocation of exported proteins across the plasma membranes must be found for both species of bacteria.

In previous studies, use of random translational gene fusions (PhoA mutagenesis) to identify and alter exported proteins in *Streptococcus pneumoniae* provided insight into putative exported proteins [Pearce et al., *Mol. Microbiol.*, 9:1037 (1993); International Patent Publication No. WO 95/06732, published March 9, 1995; U.S. Patent Application Serial No. 08/116,541, filed September 1, 1993; U.S. Patent Application Serial No. 08/245,511, filed May 18, 1994]. Coupling this gene fusion technology to bioactivity assays for pneumococcal adherence, the primary goal was to genetically identify and characterize immunogenic adhesion virulence determinants to eucaryotic cells that define the bacteria-host relationship and thus serve as vaccine candidates. Over 25 loci that effect adherence have been identified as determinants of virulence.

In addition, the molecular mechanism of pathogenesis caused by pneumococcus are beginning to be defined [Cundell, et al., *Infect. Immun.* 63:2493-2498 (1995);

Wizemann, et al., *Proc. Natl. Acad. Sci. USA* (1996); Cundell, et al., *J. Cell Biol.* **S18A**:45 (1994); Spellerberg, et al., *Mol. Microb.* (1996)]. The results of these efforts shows that many bacterial components participate in a complex coordinated process to cause disease. However, it is also apparent that this strategy has

5 produced only a few potential vaccine candidates.

Of note in the search for exported pneumococcal proteins that might be attractive targets for a vaccine is pneumococcal surface protein A (PspA) [see Yother et al., *J. Bacteriol.*, **174**:610 (1992)]. PspA has been reported to be a candidate for a *S. pneumoniae* vaccine as it has been found in all pneumococci to date; the purified protein can be used to elicit protective immunity in mice; and antibodies against the protein confer passive immunity in mice [Talkington et al., *Microb. Pathog.* **13**:343-355 (1992)]. However, PspA demonstrates antigenic variability between strains in the N-terminal half of the protein, which contains the immunogenic and

10 protection eliciting epitopes (Yother et al., *supra*). This protein does not represent a common antigen for all strains of *S. pneumoniae*, and therefore is not an optimal vaccine candidate.

15

Pneumococcal Choline Binding Proteins

20 Previous studies have shown that PspA, as well as one other surface exposed protein, LytA, the autolytic amidase, bind to teichoic acid (TA), an integral part of the cell wall of *Streptococcus pneumoniae* in a choline-dependent manner. TA contains a unique terminal phosphorylcholine moiety. PspA, a protein having a molecular weight of 84 kDa, and which is highly variable, is released from the cell

25 surface with high choline concentration. Lyt, or autolysin, is a 36 kDa protein, which lyses the pneumococcal cell wall (self lysis), but is not released from the cell by growth in high concentrations of choline, by washing in 10% choline, or by growth in ethanolamine. Reports on choline binding proteins include those by Sanchez-Puelles et al *Gene* **89**:69-75 (1990), Briese and Hakenback *Eur. J.*

30 *Biochem.* **146**:417-427, Yother and White *J. of Bacteriol.* **176**:2976-2985, Sanchez-

Beato et al *J. of Bacteriol.* **177**:1098-1103, and Wren *Micro. Review Mol. Microbiol.* **5**:797-803 (1991), which are hereby incorporated by reference in their entirety.

- 5 A variety of covalent and non-covalent mechanisms of attachment have been described for proteins decorating the surfaces of gram positive bacteria. Some streptococci and *Clostridium sp.* have phosphorylcholine as a unique component of the cell wall. This molecule is the terminal constituent of the teichoic acid (C polysaccharide) and lipoteichoic acid (LTA) attached to the cell wall and plasma
- 10 membrane of these bacteria. A family of choline binding proteins (CPBs) have also been described which serve a variety of cellular functions. These proteins consist of an N-terminal activity domain and a repeated C-terminal signature choline binding domain that anchors these molecules to the surface of the bacteria. This motif has been identified in the C-terminal regions of a secreted glycoprotein from
- 15 *Clostridium acetobutylicum* NCIB 88052 [Sanchez-Beato, et al., *J. Bacteriol.* **177**:1098-1103 (1995)], toxins A and B from *Clostridium difficile* [Von Eichel-Streiber and Sauerborn, *Gene* **96**:107-13 (1990); Von Eichel-Streiber et al., *J. Bacteriol.* **174**:6707-6710 (1992)], a glucan-binding protein from *Streptococcus mutans*, several glycosyltransferases from *Streptococcus downei* and *S. mutans*, the
- 20 murein hydrolase (LytA) from pneumococcus and pneumococcal lytic phage [Ronda et al., *Eur. J. Biochem.* **164**:621-4 (1987); Diaz et al., *J. Bacteriol.* **174**:5516-25 (1992); Romero et al., *Microb. Lett.* **108**:87-92 (1993); Yother and White, *J. Bacteriol.* **176**:2976-85 (1994)], and a surface antigen (PspA) also from pneumococcus.

25

Pathology of Pneumococcal Adherence

- S. pneumoniae* is a gram positive bacteria which is a major cause of invasive infections such as sepsis and meningitis [Tuomanen et al., *N. Engl. J. Med.* **322**:1280-1284(1995)]. The pneumococcus colonizes the nasopharyngeal
- 30 epithelium and then penetrates the epithelium of the lung or nasopharynx in order to

reach the vascular compartment. Such translocation would involve, of necessity, passage from an epithelial site through the underlying basement membrane/extracellular matrix and across endothelia. Pneumococci have been demonstrated to adhere to epithelia, endothelia and basement membrane in vitro and
 5 in vivo [Plotkowski *et al.*, *Am. Rev. Respir. Dis.*, 134 (1986); Cundell and Tuomanen, *Microb Path.*, 17:361-374 (1994); Cundel *et al.*, *Nature*, 377:435-438 (1995); van der Flier *et al.*, *Infect. Immun.*, 63:4317-4322 (1995).

Fibronectin is a mammalian glycoprotein present as a soluble dimer (molecular
 10 weight of 550 kDa) in body fluids such as plasma (200-700mg/ml), cerebrospinal fluid and amniotic fluid and as a less soluble multimer in the extracellular matrix and basement membrane [Ruoslahti, *Ann. Rev. Biochem.*, 57:375-413 (1988)]. Fibronectin has specific binding sites for a number of proteins including collagen, integrins, and two binding sites for heparin. Many microorganisms bind
 15 fibronectin, including oral streptococci and some gram negative bacteria [Westerlund and Korhonen, *Mol. Microbiol.*, 9:687-694 (1993)]. These diverse pathogens all target the Type 1 repeats of the N-terminal heparin binding domain of fibronectin. The cognate fibronectin binding proteins demonstrate a similar amino acid sequence motif consistent with binding to the same target within fibronectin
 20 [Westerlund and Korhonen, 1993, *supra*]. In contrast to this pattern, *Streptococcus pneumoniae* was found to adhere avidly to immobilized fibronectin at the carboxyterminal heparin binding domain [van der Flier *et al.*, *Infect. Immun.*, 63:4317-4322, (1995)]. This domain of fibronectin has a number of biological activities. It contains the major proteoglycan binding domain (Hep II) and also
 25 supports binding of the leukocyte integrin VLA-4 at two regions in the type III connecting segment (IIICS) [Wayner *et al.*, *Cell. Biol.*, 109:1321-1330 (1989; Guan and Hynes, *Cell*, 60:53-61 (1990); Mould *et al.*, *J. Biol. Chem.*, 266:3579-3585 (1991)]. The VLA-4 binding domain is distinct from that containing the RGD motif for binding fibronectin by VLA-5 [Pierschbacher *et al.*, *Cell*, 26:259-267 (1981).

The IIICS segment is subject to alternative splicing and is absent in some soluble forms of fibronectin [Guan and Hynes, 1990, *supra*].

VLA-4, $\alpha 4\beta 1$ (CD49d/CD29), is an integrin present on lymphocytes, monocytes,
 5 muscle cells and melanoma cells which mediates binding to VCAM-1 on endothelial cells and myoblasts and to the IIICS domain of fibronectin in the subendothelial matrix [Osborn *et al.*, *Cell*, **59**:1203-1211 (1989); Mould *et al.*, *J. Biol. Chem.* **254**:4020-4024 (1990); Shimizu *et al.*, *Immunological Reviews*, **114**:109-143 (1990); Rosen *et al.*, *Cell*, **69**:1107-1119 (1992)]. These interactions are important
 10 during infiltration of mononuclear cells to sites of inflammation, metastasis of melanoma cells and in myogenesis [McCarthy *et al.*, *J. Cell. Biol.* **102**:179-188 (1986); Osborn *et al.*, 1988, *supra*; Shimizu *et al.*, 1990, *supra*; Rosen *et al.*, 1992, *supra*]. VLA-4 targets a 25 amino acid region (CS1) with the IIICS domain of fibronectin, an interaction which can be blocked by the tripeptide Leu-Asp-Val
 15 [Guan and Hynes, 1990, *supra*; Komoriya *et al.*, *J. Biol. Chem.* **266**:15075-15079 (1991); Mould *et al.*, 1991, *supra*]. An homologous motif IDSP is present in the VLA-4 binding sites in VCAM-1 domains I and IV [Clements *et al.*, *J. Cell. Sci.*, **107**:2127-2135 (1994)]. The binding sites on VLA-4 for VCAM-1 and fibronectin have been suggested to be distinct but overlapping [Elices *et al.*, *Cell*, **50**:577-584
 20 (1990); Pulido *et al.*, *J. Biol. Chem.*, **266**:10241-10245 (1991); Vonderheide and Springer, *J. Exp. Med.*, **175**:1433-1442 (1992); Makarewicz, *J. Biol. Chem.* **269**:4005-4011 (1994)]. The ability of pneumococci to target the HepII region of fibronectin raised the possibility that these bacteria recognized the region common between HepII and VCAM-1 and that the binding was mediated by a bacterial
 25 version of VLA-4. Such interactions could promote passage of pneumococci across the basement membrane. The importance of VLA-4-VCAM-1 interactions for leukocyte trafficking to brain also suggested a role for pneumococcal transmigration into the central nervous system in meningitis.

Therefore, in view of the aforementioned deficiencies attendant with prior art methods of vaccinating against bacterial pathogens, it should be apparent that there still exists a need in the art for identifying protein antigens suitable for use as subunit vaccines, and for use in inducing antibodies suitable for use in passive immunization.

The citation of any reference herein should not be construed as an admission that such reference is available as prior art to the invention.

10 SUMMARY OF THE INVENTION

In accordance with the present invention, bacterial surface antigens are provided which are suitable for use in immunizing animals against bacterial infection. More particularly, novel choline binding proteins from streptococci, preferably pneumococci, are provided.

In a further embodiment, a method is provided for isolating and identifying choline binding proteins, and the genes encoding them.

20 In its broadest aspect, the present invention extends to streptococcal surface antigens, generally referred to herein as choline binding proteins, having the following characteristics:

- a) binding to choline; and
- b) eluting from a choline affinity chromatographic column in the presence of 10%, preferably at least 10%, choline in Dubelcco's phosphate buffered saline (DPBS);

with the proviso that the bacterial surface antigen of the present invention is not PspA or autolysin (LytA). In a preferred aspect, the choline binding protein of the invention has one or more of a characteristic selected from the group consisting of:

- 30 c) inhibiting adherence of the bacteria to host cells;

- d) being reactive with sera from patients infected or recovering from infection with the bacteria;
- e) being reactive with rabbit antisera generated against purified choline binding proteins isolated from a choline affinity column by elution in 10% choline, DPBS; and
- f) labeled by fluorescein isothiocyanate (FITC) without requiring bacterial lysis (*i.e.*, in intact bacteria).

In a specific example, the bacterial surface antigen is isolated from pneumococcus.

In a still further aspect, the present invention extends to vaccines based on the choline binding proteins.

In a particular embodiment, the present invention relates to all members of the herein disclosed family of bacterial surface antigens which bind choline, with the proviso that this group does not include PspA or LytA.

In a preferred embodiment, the invention provides a choline binding protein with a high degree of sequence similarity to enolase, particularly *B. subtilis* enolase.

The present invention also relates to an isolated nucleic acids, such as recombinant DNA molecule or cloned gene, or a degenerate variant thereof, which encodes a bacterial choline binding protein (CBP) of the invention. Preferably, the nucleic acid molecule, in particular a recombinant DNA molecule or cloned gene, encoding the CBP has a nucleotide sequence or is complementary to a DNA sequence or fragment thereof which codes on expression for an amino acid having a sequence as follows:

CBP112 XENEGSTQAATSSNMAKTEHRKAAKQVVDE

(SEQ ID NO:1)

CBP90 AREFSLEKTR (SEQ ID NO:2)

CBP84 XREFSLEKTRNIGIMAHVDAGKT (SEQ ID NO:3)

CBP80 KXXXWQXKQYLKEDGSQAANEXVFDTA

(SEQ ID NO:4)

CBP78 QKIIGIDLGTNSAVAVLEGTESKIIANPE

5 (SEQ ID NO:5)

CBP70 XXXEVAKXSQDTTAS (SEQ ID NO:6)

CBP60 XNERVKIVATLGPAVEGRG (SEQ ID NO:7)

CBP50 XIIXXVYAREVLDSRGNP (SEQ ID NO:8)

10 CBP112-Int1 EDRRNYHPTNTYK (SEQ ID NO:9)

CBP112-Int2 XDDQQAEDYA (SEQ ID NO:10)

Almost Full Length CBP50 (SEQ ID NO:19)

15 Partial CBP112 (SEQ ID NO:21)

In a specific embodiment a nucleic acid of the invention encodes at least a portion (an internal fragment) of the choline binding protein CBP50, preferably having a nucleotide sequence depicted in SEQ ID NO:18; or a portion of the 5' region of the

20 gene encoding binding protein CBP112, preferably having a nucleotide sequence depicted in SEQ ID NO:20.

The DNA sequences encoding the CBPs of the present invention, or portions thereof, may be prepared as probes to screen for complementary sequences and

25 genomic clones in the same or alternate species. The present invention extends to probes so prepared that may be provided for screening. For example, the probes may be prepared with a variety of known vectors, such as the phage λ vector. Such probes are useful for diagnosis, *e.g.*, to confirm a species or strain of Gram positive bacterial infection, as well as for cloning cDNA or genes encoding CBPs.

30

The present invention also includes CBPs having the activities noted herein, and that display the amino acid sequences set forth and described above and selected from SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 19, and 21.

- 5 In a further embodiment of the invention, the full DNA sequence of the recombinant DNA molecule or cloned gene so determined may be operatively linked to an expression control sequence which may be introduced into an appropriate host. The invention accordingly extends to unicellular hosts transformed with the cloned gene or recombinant DNA molecule comprising a
- 10 DNA sequence encoding the present CBP(s), and more particularly, the DNA sequences or fragments thereof determined from the sequences set forth above and in SEQ ID NOS:18 and 20.

- According to other preferred features of certain preferred embodiments of the
- 15 present invention, a recombinant expression system is provided to produce biologically active CBPs or immunologically reactive portions thereof.

- The concept of the bacterial surface antigens contemplates that specific factors exist for correspondingly specific binding proteins, such as CBPs and the like, as
- 20 described earlier. Accordingly, the exact structure of each CBP will understandably vary so as to achieve this choline-binding and activity specificity. It is this specificity and the direct involvement of the CBP in the adherence of the bacteria, that offers the promise of a broad spectrum of diagnostic and therapeutic utilities.

- 25 The present invention naturally contemplates several means for preparation of the CBPs, including as illustrated herein known recombinant techniques, and the invention is accordingly intended to cover such synthetic preparations within its scope. The isolation of the DNA and amino acid sequences disclosed herein facilitates the reproduction of the CBPs by such recombinant techniques, and
- 30 accordingly, the invention extends to expression vectors prepared from the disclosed

DNA sequences for expression in host systems by recombinant DNA techniques, and to the resulting transformed hosts.

- A particular advantage of the present invention is that it provides for preparation of
- 5 sufficient quantities of CBPs for commercialization of anti-CBP vaccines. A further advantage is that the invention provides for preparation of multi-component vaccines containing two or more CBPs, thus broadening and increasing the potential effectiveness of the vaccine. In its primary aspect, the invention contemplates utilizing the CBPs of the invention, either separately or in combinations of two or
- 10 more, in vaccines for protection against pneumococcal infection. Preferably, a vaccine comprising two or more CBPs further comprises PspA, LytA, or both. According to the invention, CBPs may be prepared recombinantly. Alternatively, CBPs may be obtained from bacterial cultures, *e.g.*, using the purification methods described and exemplified herein. In a specific embodiment, a mixture of CBPs
- 15 from pneumococcus are obtained, *e.g.*, by choline affinity chromatography, and are used directly without further purification to immunize an animal and elicit protective antibodies. Such a mixture of choline affinity purified proteins may be obtained from bacteria that express PspA or that lack PspA expression (*i.e.*, PspA⁻ bacteria as exemplified herein).
- 20 In another aspect, the genes (*e.g.*, cDNA) encoding one or more CBPs of the invention are engineered in a transgenic vector for expression in a mammalia host *in vitro*, as a nucleic acid-based vaccine.
- 25 In yet another embodiment, the CBPs of the invention are used to generate antibodies for passive immunization, diagnostics, or screening. In a specific example, *infra*, passive immunization prevents death from pneumococcal infection in a murine model.

The diagnostic utility of the present invention extends to the use of binding partners, notably antibodies, to the present CBPs in assays to screen for bacterial infection.

- 5 Antibodies against the CBP(s) include naturally raised and recombinantly prepared antibodies. Such antibodies could be used to screen expression libraries to obtain the gene or genes that encode the CBP(s). These may include both polyclonal and monoclonal antibodies prepared by known genetic techniques, as well as bi-specific (chimeric) antibodies, and antibodies including other functionalities suiting them for
10 additional diagnostic use conjunctive with their capability of modulating bacterial adherence.

- Thus, the CBPs, their analogs and/or analogs, and antibodies that may be raised thereto, are capable of use in connection with various diagnostic techniques,
15 including immunoassays, such as a radioimmunoassay, using for example, an antibody to the CBP that has been labeled by either radioactive addition, or radioiodination.

- In an immunoassay, a control quantity of the antagonists or antibodies thereto, or
20 the like may be prepared and labeled with an enzyme, a specific binding partner and/or a radioactive element, and may then be introduced into a cellular sample. After the labeled material or its binding partner(s) has had an opportunity to react with sites within the sample, the resulting mass may be examined by known techniques, which may vary with the nature of the label attached.

25

In the instance where a radioactive label, such as the isotopes ^3H , ^{14}C , ^{32}P , ^{35}S , ^{36}Cl , ^{51}Cr , ^{57}Co , ^{58}Co , ^{59}Fe , ^{90}Y , ^{125}I , ^{131}I , and ^{186}Re are used, known currently available counting procedures may be utilized. In the instance where the label is an enzyme, detection may be accomplished by any of the presently utilized colorimetric,

spectrophotometric, fluorospectrophotometric, amperometric or gasometric techniques known in the art.

The present invention includes an assay system which may be prepared in the form
 5 of a test kit for the quantitative analysis of the extent of the presence of the bacterial
 infection, or to identify drugs or other agents that may mimic or block such
 infection. The system or test kit may comprise a labeled component prepared by
 one of the radioactive and/or enzymatic techniques discussed herein, coupling a
 label to the CBPs, their agonists and/or antagonists, and one or more additional
 10 immunochemical reagents, at least one of which is a free or immobilized ligand,
 capable either of binding with the labeled component, its binding partner, one of the
 components to be determined or their binding partner(s).

In particular, the proteins of CBPs whose sequences are presented in SEQ ID
 15 NOS:1-10, 19 and 20 herein, their antibodies, agonists, antagonists, or active
 fragments thereof, could be prepared in pharmaceutical formulations for
 administration in instances wherein antibiotic therapy is appropriate, such as to treat
 or prevent bacterial infection. Such free proteins could compete with bacterial CBP
 function, thus interfering with bacterial pathological activity such as adhesion.

20 In a preferred aspect, the invention provides a method and associated compositions
 for treating infection with a bacterium that expresses a streptococcal choline binding
 protein comprising administering pulmonarily an adhesion inhibitory agent selected
 from the group consisting of a choline binding protein having the following

25 characteristics:

choline-binding activity; and

elution from a chromatographic column in the presence of 10% choline;

with the proviso that the streptococcal choline binding protein is not PspA or
 autolysin (LytA), an antibody to the choline binding protein, an enolase, a hindered

30 cationic small molecule, the peptide WQPPRARI (SEQ ID NO:11), and an antibody

specific for an epitope having the amino acid sequence WQPPRARI (SEQ ID NO:11). Preferably, the hindered cationic small molecule is selected from the group consisting of lysine, choline, and arginine. In a further embodiment, the adhesion inhibitory agent is administered with another drug, such as an antibiotic, a
5 steroid, a non-steroidal anti-inflammatory drug, etc.

Accordingly, it is a principal object of the present invention to provide a CBP and its subunits in purified form.

10 It is a further object of the present invention to provide antibodies to the CBP and its subunits, and methods for their preparation, including recombinant means.

It is a further object of the present invention to provide a method for detecting the presence of the CBP and its subunits in mammals in which invasive, spontaneous,
15 or idiopathic pathological states are suspected to be present.

It is a still further object of the present invention to provide method for the treatment of mammals to control the amount or activity of the bacteria, the CBP or subunits thereof, so as to alter the adverse consequences of such presence or
20 activity, or where beneficial, to enhance such activity.

It is a still further object of the present invention to provide a method for the treatment of mammals to control the amount or activity of the bacteria or its subunits, so as to treat or avert the adverse consequences of invasive, spontaneous,
25 or idiopathic pathological states.

It is a still further object of the present invention to provide pharmaceutical compositions for use in therapeutic methods which comprise or are based upon the CBP, its subunits, or their binding partner(s).

30

Other objects and advantages will become apparent to those skilled in the art from a review of the ensuing description which proceeds with reference to the following illustrative drawings and Detailed Description of the Invention.

5

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1. Western blot of a 6% SDS-PAGE on which bacterial surface proteins were separated. Lane A represents FITC-labelled surface proteins from LM91 (PspA⁻). Lane B represents FITC-labelled surface proteins from LM91 after
10 choline affinity chromatography. Lane C represents LM91 proteins after choline affinity chromatography (CBRO). "FITC" indicates anti-FITC (mouse) antibody was used. "Convalesc." indicates human convalescent (post-pneumococcal infection) antibody was used. "CBP" indicates anti-CBP antibody (rabbit) was used.

15

FIGURE 2. Western blot of a 10% SDS-PAGE as described in FIGURE 1.

FIGURE 3. Adhesion assay of PN R6 to lung cell A549.

20 **FIGURE 4.** Adhesion assay of PN R6 to HUVEC (endothelial) cells.

FIGURE 5. PRIOR ART. Schematic drawing of fibronectin and its fragments.

FIGURE 6. Direct adherence of pneumococcus to recombinant fragments of the 33
25 kDa heparin 11 binding domain.

FIGURE 7. Direct adherence of pneumococcus to synthetic peptides based on the heparin 11 type III #14 and IIICS regions.

FIGURE 8. Inhibition of *S. pneumoniae* adherence by choline. Bacteria were preincubated with choline, eluted proteins and choline were washed away, and bacterial adhesion to fibronectin evaluated.

- 5 **FIGURE 9.** Inhibition of pneumococcal adherence to fibronectin by an anti-choline antibody, TEPC15.

FIGURE 10. Comparison of amino terminal sequence of 50 kDa protein with *B. subtilis* enolase.

10

FIGURE 11. Inhibition of pneumococcal adherence to fibronectin by yeast enolase (Sigma).

FIGURE 12. Inhibition of pneumococcal adherence to fibronectin by L-lysine.

15

FIGURE 13. SDS-PAGE of eluate from a fibronectin-coupled CnBr-activated SEPHAROSE 4B column (lane A), and yeast enolase (Sigma) (lane B). French Press lysate of *S. pneumoniae* was applied to the fibronectin-SEPHAROSE column. The column was washed and adsorbed proteins eluted in 0.01, 0.1, and 0.5 M L-lysine. The 0.5 M lysine eluate was analyzed by SDS-PAGE with silver stain. The eluted protein had an apparent molecular weight comparable to yeast enolase.

20

FIGURE 14. DNA Sequence comparison of DNA for the 50 kDa protein (SEQ ID NO:14) and *B. subtilis* enolase (SEQ ID NO:15). The sequences were 74% identical.

25

FIGURE 15. Deduced amino acid sequence comparison for the 50 kDa protein (SEQ ID NO:16) and *B. subtilis* enolase (SEQ ID NO:17). These sequences were 72% identical, with 85% positives.

30

FIGURE 16. Passive protection of mice against sepsis with anti-CBP antiserum.

(A) CF1 mice challenged intraperitoneally with D39 (type 2) pneumococci at a dose of either 4.5×10^4 (n=10 per group; ●) or 8.4×10^4 (n=5 per group; ■) in 0.5 ml PBS. Experimental groups (—) were treated with 0.5 ml of 1:10 diluted rabbit anti-CBP serum one hour after inoculation with bacteria. Control mice (—) received pre-immune serum. Percent survival was evaluated daily over a six day period. (B) Passive protection against challenge with heterologous serotype SIII (type 3; n=5 per group; □) was performed as in A with a dose of 200 cfu. Bacteria were preincubated with antiserum (●) for 30 min prior to intraperitoneal inoculation. Blood was sampled at 24 hours for cfu and survival was assessed every 12 hours for 72 hours.

FIGURE 17. Competitive inhibition of pneumococcal adherence to human cells by exogenous purified CBPs. (A) Type II lung cells (LC) or (B) endothelial cells (EC)

were preincubated for 15 min with CBP preparation (1 mg/ml), washed and then incubated with FITC-labelled R6 pneumococcus. *E. coli* DH5a was used as a control. Adherence was quantitated by fluorescence intensity. (C) Dose dependence of the ability of CBPs to inhibit adherence of FITC-labelled pneumococci.

FIGURE 18. Adherence of CbpA⁻ mutant to human cells and glycoconjugates.

(A) Monolayers of Type II lung cell line A549 (LC) or (B) umbilical vein endothelial cells (EC) were activated by treatment with TNF α (10 ng/ml for 2 h) or IL-1 (5 ng/ml for 4 h) or were untreated (resting). R6 and the CbpA⁻ mutant were fluorescein labelled and overlaid onto the monolayers for 30 min. Nonadherent bacteria were washed away and adherence was quantitated visually and expressed as a % of control, i.e. 100% = 498 ± 81 or 174 ± 81 R6 bacteria per 100 resting LC and EC respectively. The experiment was performed 5 times with 6-8 wells per condition. (C) Terasaki plates were coated with 6'sialyllactose (6'SL), lacto-N-neotetraose (LnNT), N-acetylgalactosamine- β 1,4-galactose HSA, (GlcNAc-

5 β 1,4-Gal), N-acetylgalactosamine- β 1,3-galactose HSA (GlcNAc β 1,3-Gal), or HSA at 100 μ M. R6 (solid bars) or CbpA⁻ (stippled bars) were grown to an OD₆₂₀ of 0.4 for all experiments except those with 6'SL (OD₆₂₀ of 0.6). Fluorescein labelled bacteria were placed in the wells for 30 min, the wells were washed, and adherent
 10 bacteria were counted visually per 40x field. Values are expressed as a % of control, i.e. 100% = 133 \pm 24 R6 per 40 x field of 6'SL well, 32 \pm 8, 20 \pm , and 15 \pm 4 for 6'SL, LnNT, GlcNAc- β 1,4-gal, and GlcNAc β 1,3-Gal, respectively. The experiment was performed 3 times with 6-12 wells per condition.

10 **FIGURE 19.** The contribution of CbpA to pneumococcal carriage in infant rats. The ability to colonize the nasopharynx of 5 day old rats were compared between the control strains D39 (hatched), D39 (iga⁻) (stippled) and an isogenic *cbpA* deficient strain (solid). The ordinate indicates the time following inoculation and the abscissa the number of colonies in nasal washes (mean \pm SD, n=20 per group).

15

DETAILED DESCRIPTION

As noted above, the present invention is directed to bacterial surface antigens that are suitable for use in immunizing animals against bacterial infection. More
 20 particularly, novel choline binding proteins from pneumococci are provided. These proteins, which are found at the surface of pneumococci, when formulated with an appropriate adjuvant, are used in vaccines for protection against pneumococci, and against other bacteria with cross-reactive choline binding proteins.

25 As has been previously reported, *S. pneumoniae* adheres to fibronectin at a site within the carboxy-terminal heparin II binding domain [Flier *et al.*, *Infect. Immun.* **63**:4317 (1995)]. An eight amino acid stretch within the type III # 14 repeat supports adherence. The present invention is based, in part, on the discovery that the pneumococcal adhesin for fibronectin appears to be a choline-binding protein of
 30 approximately 50 kDa. This protein has significant homology to the glycolytic

- enzyme enolase. For example, preincubation of *S. pneumoniae* with rsVCAM inhibits adherence to whole fibronectin by 96%, and *S. pneumoniae* adhere directly to rsVCAM with 6% of adherence to whole fibronectin. In addition, pneumococci bind directly to a synthetic peptide, Fn5, based on the Heparin II type III #14
- 5 region, having the sequence WQPPRARI (SEQ ID NO:11). Antibody to Fn5 inhibits adherence of *S. pneumoniae* to whole fibronectin by greater than 70% (1:100). The data exemplified herein show that *S. pneumoniae* grown in ethanolamine vs. choline results in a greater than 90% decrease in adherence. Preparations of CBP competitively inhibit adherence of *S. pneumoniae* to
- 10 fibronectin by 75-90%. Anti-choline antibody inhibits adherence of *S. pneumoniae* to fibronectin by greater than 95% at dilution of 1:10 to 1:5000. Pretreatment of *S. pneumoniae* in 10% choline results in greater than 50% decrease in adherence to fibronectin.
- 15 In a particular embodiment, the present invention relates to all members of the herein disclosed pneumococcal CBPs, with the proviso that this group does not include Lyt or PspA. In a specific embodiment, a CBP of the invention is a homolog of enolase.
- 20 The possibilities both diagnostic and therapeutic that are raised by the existence of the CBP, derive from the fact that the CBP appears to participate in direct and causal protein-protein interaction between the bacteria and its host cell. As suggested earlier and elaborated further on herein, the present invention contemplates pharmaceutical intervention in the binding reaction in which the CBP
- 25 is implicated, to modulate the activity initiated by the binding of the CBP.

In a further embodiment, a method is provided for isolating and identifying choline binding proteins, and the genes encoding them. More particularly, isolation of the genes encoding choline binding proteins of the invention allows for recombinant

production of the proteins, which greatly increases the ability to generate cost-effective vaccines.

The choline binding protein streptococcal surface antigens of the invention have the following characteristics:

- a) binding to choline; and
- b) eluting from a choline affinity chromatographic column in the presence of 10% choline in Dulbecco's phosphate buffered saline (DPBS);
- 10 with the proviso that the streptococcal surface antigen of the present invention is not PspA or autolysin (LytA). In a preferred aspect, the choline binding protein of the invention has one or more of a characteristic selected from the group consisting of:
 - c) inhibiting adherence of the bacteria to host cells;
 - d) being reactive with sera from patients infected or recovering from
 - 15 infection with the bacteria;
 - e) being reactive with rabbit antisera generated against purified choline binding proteins isolated from a choline affinity column by elution in 10% choline, DPBS; and
 - f) labeled by fluorescein isothiocyanate (FITC) without requiring
 - 20 bacterial lysis (*i.e.*, in intact bacteria).

In a specific example, *infra*, the streptococcal surface antigen is isolated from pneumococcus by affinity chromatography on a choline agarose column by elution with 10% choline.

25

In a further embodiment, exemplified herein, a peptide based on the heparin II type III #14 region of fibronectin binds the streptococcal enolase homolog of the invention. In a specific embodiment, the peptide is Fn5 having an amino acid sequence WQPPRARI (SEQ ID NO:11). Whole pneumococci adhere to this

30 peptide prepared synthetically. Thus, the free peptide would be expected to inhibit

enolase-mediated adherence to fibronectin *in vivo*. In addition, an antibody specific for this peptide inhibits pneumococcal adherence to fibronectin. In a specific embodiment, an anti Fn5 antibody inhibits adherence of *S. pneumoniae* to whole fibronectin by greater than 70%.

5

The term "bacterial" used herein refers to Gram positive bacteria with choline binding proteins homologous to the proteins exemplified herein. The present invention is more particularly directed to streptococcal CBPs and most particularly to pneumococcal CBPs.

10

The terms "bacterial (or streptococcal or pneumococcal) surface antigen", "choline binding protein (CBP)" and any variants not specifically listed, may be used herein interchangeably, and as used throughout the present application and claims refer to proteinaceous material including single or multiple proteins, and extends to those proteins having the amino acid sequence data described herein and identified by (SEQ ID NOS:1-10, 19, and 20), and the profile of activities set forth herein and in the Claims. Accordingly, proteins displaying substantially equivalent or altered activity are likewise contemplated. These modifications may be deliberate, for example, such as modifications obtained through site-directed mutagenesis, or may be accidental, such as those obtained through mutations in hosts that are producers of the complex or its named subunits. Also, the terms "bacterial (or streptococcal or pneumococcal) surface antigen", and "choline binding protein (CBP)" are intended to include within their scope proteins specifically recited herein as well as all substantially homologous analogs and allelic variations.

25

The term "enolase" refers to the enzyme 2-phospho-D-glycerate hydrolyase.

The amino acid residues described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property of immunoglobulin-

30

binding is retained by the polypeptide. NH_2 refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. Abbreviations used herein are in keeping with standard polypeptide nomenclature, *J. Biol. Chem.*, **243**:3552-59

5 (1969).

It should be noted that all amino-acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the
10 beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino-acid residues.

Purification of CBPs

15 Teichoic acid (TA), an integral part of the cell wall of *Streptococcus pneumoniae* contains a unique terminal phosphorylcholine moiety. Choline affinity chromatography or Mono-Q Sepharose, a close relative of choline were used to purify the CBPs. It is important to note that initially these proteins were purified from a capsulated strain of pneumococcus that was genetically altered not to
20 produce PspA, a major CBP. The purification schemes are as follows:

Whole bacteria are incubated with either choline-agarose or Mono-Q Sepharose.

25 Bacteria are lysed with detergent and unbound material washed with 0.5 M NaCl.

CBPs are eluted with either with 0.5 M choline chloride or with a linear choline chloride gradient. The molecular masses of proteins purified by
30 these two methods are summarized in Table 1 (see Figure 1).

Table 1: Criteria for the identification of choline binding proteins as vaccine candidates from a capsulated PspA deficient strain of pneumococcus*

Purification Scheme	SDS-PAGE 10%	Labeling of surface proteins	Rabbit antisera to whole bacteria	Human convalescent serum
Cell lysate (0.5% Triton X 100)		112 kDa 80 kDa 60 kDa 37 kDa 28 kDa	200 kDa 90 kDa 75 kDa 48 kDa 37 kDa	200 kDa 112 kDa 84 kDa 80 kDa 60 kDa
Mono-Q Anion			5% choline gradient: 112 kDa 53 kDa	
Choline-Agarose (Elution with 0.5 M choline chloride)	200 kDa 90 kDa 82 kDa 75 kDa 40 kDa	112 kDa 80 kDa 70 kDa	75 kDa < 35 kDa	112 kDa 80 kDa 75 kDa

*Bold faced characters represent proteins of interest that are potential vaccine candidates.

Purification of CBPs from the choline agarose affinity chromatography is preferred, with at least 9 proteins with molecular masses ranging from 200 to 40 kDa identifiable this way. A band of 37 kDa was also detected and shown to be LytA with antisera specific for this protein. This is a well characterized CBP and thus served as a positive control.

Criteria for Vaccine Candidates

CBPs may be subjected to a variety of tests to determine if they are good vaccine candidates. The criteria for vaccine development, and the characteristics of the isolated CBPs are summarized below.

The CBPs must be surface exposed. Whole bacteria may be chemically labeled with FITC (Fluorescein isothiocyanate) and labeled proteins detected with antisera specific for FITC. (Table 1 and Figure 1). The CBPs 112, 75, and 80, as well as LytA, were effectively labeled with FITC suggesting that these proteins were surface exposed.

Vaccine candidates must be immunogenic therefore candidate CBPs should react with human convalescent sera. The CBPs 112, 75, and 80 gave a strong signal with pooled human antisera obtained from individuals recovering from pneumococcal disease (Figure 1).

Vaccine candidates must be antigenic. The CBPs from choline agarose chromatography were injected into rabbits and sera tested for cross reactivity. Several proteins produced a strong signal. Prominent were CBPs 112, 90, 84 and 70, and 50 (Figures 1 and 2).

A good vaccine candidate may block adherence to target cell receptors present within the host at the critical sites of infection. Compared to controls, the CBP

fraction blocked pneumococcal adherence by 45% to LC and 89% to EC in a dose dependent manner (Figure 3). Based on these results it is most likely that the fraction of CBPs contain adhesins involved in the binding of bacteria to eucaryotic target cells. The contribution of each of these CBPs to block the adhesive

- 5 properties of parental bacteria can be assessed, for example, to block the binding of pneumococcus to epithelial (type II lung cells), endothelial (human umbilical vein endothelial cells) cells, and immobilized glycoconjugates that contain GlcNAc β 1-4Gal, GlcNAc β 1-3Gal, GlcNAc, or other sugars that have been shown to be analogs for eucaryotic receptors.

10

It should be noted that not every CBP may function as an adhesin, similarly, adhesion activity may be a collateral characteristic of CBPs.

- 15 A preferred vaccine candidate will elicit a protective immune response without antigenic variability among in clinical serotypes. Antisera (either monoclonal or polyclonal) to each CBP will be generated to determine if the native and the recombinant CBPs are immunogenic. CBP specific antibodies will be used to screen relevant pneumococcal serotypes for antigenic variability.

- 20 In a more preferred aspect, antibodies to the CBPs will be tested to confirm that they protect against pneumococcal infection, preferably of various strains or serotypes. For example, passively or actively immunized animals are challenged with pneumococcus in models for bacteremia or colonization, or preferably both.

- 25 Other criteria to consider in selection of a preferred CBP as a vaccine candidate include testing CBP defective mutants for attenuation of virulence in animal models for bacteremia or colonization efficacy alone or in combination or coupled to a capsular polysaccharide. For example, preliminary data show that CBP112 is expressed in virulent transparent bacteria, but that expression is diminished in
30 avirulent opaque bacteria.

Genes Encoding CBPs

As stated above, the present invention also relates to a recombinant DNA molecule or cloned gene, or a degenerate variant thereof, which encodes a CBP, or a
 5 fragment thereof, that possesses a molecular weight of between about 50 kDa and 112 kDa, preferably which has an amino acid sequence set forth in (SEQ ID NOS:1-10, 19 and 20). In a specific aspect, a nucleic acid molecule, in particular a recombinant DNA molecule or cloned gene, encoding the 50-112 kDa CBP has a nucleotide sequence or is complementary to a DNA sequence shown in (SEQ ID
 10 NO:18 or 20).

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g.,
 15 Sambrook et al, "Molecular Cloning: A Laboratory Manual" (1989); "Current Protocols in Molecular Biology" Volumes I-III [Ausubel, R. M., ed. (1994)]; "Cell Biology: A Laboratory Handbook" Volumes I-III [J. E. Celis, ed. (1994)]; "Current Protocols in Immunology" Volumes I-III [Coligan, J. E., ed. (1994)]; "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization"
 20 [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription And Translation" [B.D. Hames & S.J. Higgins, eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

25 Therefore, if appearing herein, the following terms shall have the definitions set out below.

A "replicon" is any genetic element (e.g., plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*; i.e., capable of
 30 replication under its own control.

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

- 5 A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in its either single stranded form, or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear DNA molecules (e.g.,
- 10 restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (i.e., the strand having a sequence homologous to the mRNA).

15

An "origin of replication" refers to those DNA sequences that participate in DNA synthesis.

- A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed
- 20 and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences
- 25 from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. A polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for the expression of a coding sequence in a host cell.

- 5 A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements
- 10 necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Prokaryotic
- 15 promoters contain Shine-Dalgarno sequences in addition to the -10 and -35 consensus sequences.

- An "expression control sequence" is a DNA sequence that controls and regulates the transcription and translation of another DNA sequence. A coding sequence is
- 20 "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then translated into the protein encoded by the coding sequence.

- A "signal sequence" can be included before the coding sequence. This sequence
- 25 encodes a signal peptide, N-terminal to the polypeptide, that communicates to the host cell to direct the polypeptide to the cell surface or secrete the polypeptide into the media, and this signal peptide is clipped off by the host cell before the protein leaves the cell. Signal sequences can be found associated with a variety of proteins native to prokaryotes and eukaryotes.

The term "oligonucleotide," as used herein in referring to the probe of the present invention, is defined as a molecule comprised of two or more ribonucleotides, preferably more than three. Its exact size will depend upon many factors which, in turn, depend upon the ultimate function and use of the oligonucleotide.

5

- The term "primer" as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product, which is complementary to a
- 10 nucleic acid strand, is induced, i.e., in the presence of nucleotides and an inducing agent such as a DNA polymerase and at a suitable temperature and pH. The primer may be either single-stranded or double-stranded and must be sufficiently long to prime the synthesis of the desired extension product in the presence of the inducing agent. The exact length of the primer will depend upon many factors, including
- 15 temperature, source of primer and use of the method. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides.
- 20 A DNA sequence is "operatively linked" to an expression control sequence when the expression control sequence controls and regulates the transcription and translation of that DNA sequence. The term "operatively linked" includes having an appropriate start signal (e.g., ATG) in front of the DNA sequence to be expressed and maintaining the correct reading frame to permit expression of the
- 25 DNA sequence under the control of the expression control sequence and production of the desired product encoded by the DNA sequence. If a gene that one desires to insert into a recombinant DNA molecule does not contain an appropriate start signal, such a start signal can be inserted in front of the gene.

The term "standard hybridization conditions" refers to salt and temperature conditions substantially equivalent to 5 x SSC and 65°C for both hybridization and wash. However, one skilled in the art will appreciate that such "standard hybridization conditions" are dependent on particular conditions including the

5 concentration of sodium and magnesium in the buffer, nucleotide sequence length and concentration, percent mismatch, percent formamide, and the like. Also important in the determination of "standard hybridization conditions" is whether the two sequences hybridizing are RNA-RNA, DNA-DNA or RNA-DNA. Such standard hybridization conditions are easily determined by one skilled in the art

10 according to well known formulae, wherein hybridization is typically 10-20°C below the predicted or determined T_m with washes of higher stringency, if desired.

The primers herein are selected to be "substantially" complementary to different strands of a particular target DNA sequence. This means that the primers must be

15 sufficiently complementary to hybridize with their respective strands. Therefore, the primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the strand. Alternatively, non-complementary bases or longer sequences can be

20 interspersed into the primer, provided that the primer sequence has sufficient complementarity with the sequence of the strand to hybridize therewith and thereby form the template for the synthesis of the extension product.

As used herein, the terms "restriction endonucleases" and "restriction enzymes"

25 refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be

30 integrated (covalently linked) into chromosomal DNA making up the genome of the

cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells

5 through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

10

Two DNA sequences are "substantially homologous" when at least about 75% (preferably at least about 80%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using

15 standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Maniatis et al., *supra*; DNA Cloning, Vols. I & II, *supra*; Nucleic Acid Hybridization, *supra*.

20

It should be appreciated that also within the scope of the present invention are DNA sequences encoding CBPs which code for a CBP having the same amino acid sequence as one of SEQ ID NOS:1-5, but which are degenerate to an oligonucleotide represented by the purified natural (*i.e.*, native) gene, *e.g.*, by SEQ

25 ID NO:11. By "degenerate to" is meant that a different three-letter codon is used to specify a particular amino acid. It is well known in the art that the following codons can be used interchangeably to code for each specific amino acid:

Phenylalanine (Phe or F)	UUU or UUC
30 Leucine (Leu or L)	UUA or UUG or CUU or CUC or CUA or CUG

	Isoleucine (Ile or I)	AUU or AUC or AUA
	Methionine (Met or M)	AUG
	Valine (Val or V)	GUU or GUC or GUA or GUG
	Serine (Ser or S)	UCU or UCC or UCA or UCG or AGU or AGC
5	Proline (Pro or P)	CCU or CCC or CCA or CCG
	Threonine (Thr or T)	ACU or ACC or ACA or ACG
	Alanine (Ala or A)	GCU or GCG or GCA or GCC
	Tyrosine (Tyr or Y)	UAU or UAC
	Histidine (His or H)	CAU or CAC
10	Glutamine (Gln or Q)	CAA or CAG
	Asparagine (Asn or N)	AAU or AAC
	Lysine (Lys or K)	AAA or AAG
	Aspartic Acid (Asp or D)	GAU or GAC
	Glutamic Acid (Glu or E)	GAA or GAG
15	Cysteine (Cys or C)	UGU or UGC
	Arginine (Arg or R)	CGU or CGC or CGA or CGG or AGA or AGG
	Glycine (Gly or G)	GGU or GGC or GGA or GGG
	Termination codon	UAA (ochre) or UAG (amber) or UGA (opal)

- 20 It should be understood that the codons specified above are for RNA sequences. The corresponding codons for DNA have a T substituted for U.

Completely degenerate oligonucleotides can be designed based on the amino acid sequence encoding the CBP, taking into account the above degenerates. Likewise, 25 where a codon encoding an amino acid is not known with certainty, *e.g.*, due to degeneracy of the genetic code, inosine may be included at the unknown position.

Mutations can be made in a nucleic acid encoding a CBP such that a particular codon is changed to a codon which codes for a different amino acid. Such a 30 mutation is generally made by making the fewest nucleotide changes possible. A

substitution mutation of this sort can be made to change an amino acid in the resulting protein in a non-conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to another grouping) or in a conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to the same grouping). Such a conservative change generally leads to less change in the structure and function of the resulting protein. A non-conservative change is more likely to alter the structure, activity or function of the resulting protein. The present invention should be considered to include sequences containing conservative changes which do not significantly alter the activity or binding characteristics of the resulting protein. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point.

Particularly preferred substitutions are:

- Lys for Arg and vice versa such that a positive charge may be maintained;
- Glu for Asp and vice versa such that a negative charge may be maintained;
- Ser for Thr such that a free -OH can be maintained; and
- Gln for Asn such that a free NH_2 can be maintained.

Two amino acid sequences are "substantially homologous" when at least about 70% of the amino acid residues (preferably at least about 80%, and most preferably at least about 90 or 95%) are identical, or represent conservative substitutions.

- 5 A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. Another example of a
- 10 heterologous coding sequence is a construct where the coding sequence itself is not found in nature (e.g., a cDNA where the genomic coding sequence contains introns, or synthetic sequences having codons different than the native gene). Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

15

The present invention extends to the preparation of oligonucleotides that may be used to hybridize with nucleic acids encoding CBPs.

- As mentioned above, a DNA sequence encoding a CBP can be prepared
- 20 synthetically rather than cloned. The DNA sequence can be designed with the appropriate codons for the CBP amino acid sequence. In general, one will select preferred codons for the intended host if the sequence will be used for expression. The complete sequence is assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence. See, e.g., Edge,
- 25 *Nature*, **292**:756 (1981); Nambair et al., *Science*, **223**:1299 (1984); Jay et al., *J. Biol. Chem.*, **259**:6311 (1984).

Synthetic DNA sequences allow convenient construction of genes which will express CBP analogs or "muteins". Alternatively, DNA encoding muteins can be

made by site-directed mutagenesis of native CBP genes or cDNAs, and muteins can be made directly using conventional polypeptide synthesis.

- 5 A general method for site-specific incorporation of unnatural amino acids into proteins is described in Christopher J. Noren, Spencer J. Anthony-Cahill, Michael C. Griffith, Peter G. Schultz, *Science*, **244**:182-188 (April 1989). This method may be used to create analogs with unnatural amino acids.

Recombinant Production of CBPs

- 10 Another feature of this invention is the expression of the DNA sequences disclosed herein. As is well known in the art, DNA sequences may be expressed by operatively linking them to an expression control sequence in an appropriate expression vector and employing that expression vector to transform an appropriate unicellular host.
- 15 Such operative linking of a DNA sequence of this invention to an expression control sequence, of course, includes, if not already part of the DNA sequence, the provision of an initiation codon, ATG, in the correct reading frame upstream of the DNA sequence.
- 20 A wide variety of host/expression vector combinations may be employed in expressing the DNA sequences of this invention. Useful expression vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Suitable vectors include derivatives of SV40 and known bacterial plasmids, e.g., *E. coli* plasmids col El, pCR1, pBR322, pMB9 and their
- 25 derivatives, plasmids such as RP4; phage DNAs, e.g., the numerous derivatives of phage λ , e.g., NM989, and other phage DNA, e.g., M13 and filamentous single stranded phage DNA; yeast plasmids such as the 2μ plasmid or derivatives thereof; vectors useful in eukaryotic cells, such as vectors useful in insect or mammalian
- 30 cells; vectors derived from combinations of plasmids and phage DNAs, such as

plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

Any of a wide variety of expression control sequences -- sequences that control the expression of a DNA sequence operatively linked to it -- may be used in these vectors to express the DNA sequences of this invention. Such useful expression control sequences include, for example, the early or late promoters of SV40, CMV, vaccinia, polyoma or adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the *LTR* system, the major operator and promoter regions of phage λ , the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase (e.g., Pho5), the promoters of the yeast α -mating factors, and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof.

A wide variety of unicellular host cells are also useful in expressing the DNA sequences of this invention. These hosts may include well known eukaryotic and prokaryotic hosts, such as strains of *E. coli*, *Pseudomonas*, *Bacillus*, *Streptomyces*, fungi such as yeasts, and animal cells, such as CHO, R1.1, B-W and L-M cells, African Green Monkey kidney cells (e.g., COS 1, COS 7, BSC1, BSC40, and BMT10), insect cells (e.g., Sf9), and human cells and plant cells in tissue culture.

It will be understood that not all vectors, expression control sequences and hosts will function equally well to express the DNA sequences of this invention. Neither will all hosts function equally well with the same expression system. However, one skilled in the art will be able to select the proper vectors, expression control sequences, and hosts without undue experimentation to accomplish the desired expression without departing from the scope of this invention. For example, in selecting a vector, the host must be considered because the vector must function in it. The vector's copy number, the ability to control that copy number, and the

expression of any other proteins encoded by the vector, such as antibiotic markers, will also be considered.

In selecting an expression control sequence, a variety of factors will normally be considered. These include, for example, the relative strength of the system, its controllability, and its compatibility with the particular DNA sequence or gene to be expressed, particularly as regards potential secondary structures. Suitable unicellular hosts will be selected by consideration of, e.g., their compatibility with the chosen vector, their secretion characteristics, their ability to fold proteins correctly, and their fermentation requirements, as well as the toxicity to the host of the product encoded by the DNA sequences to be expressed, and the ease of purification of the expression products.

Considering these and other factors a person skilled in the art will be able to construct a variety of vector/expression control sequence/host combinations that will express the DNA sequences of this invention on fermentation or in large scale animal culture.

It is further intended that CBP analogs may be prepared from nucleotide sequences of the protein complex/subunit derived within the scope of the present invention. Analogs, such as fragments, may be produced, for example, by pepsin digestion of CBP or bacterial material. Other analogs, such as muteins, can be produced by standard site-directed mutagenesis of CBP coding sequences. Analogs exhibiting "choline-binding activity" such as small molecules, whether functioning as promoters or inhibitors, may be identified by known *in vivo* and/or *in vitro* assays.

Antibodies to CBPs

As noted above, the CBPs of the invention, whether obtained by purification from bacterial sources or recombinantly, can be used to generate antibodies for diagnosis

and therapy, as set forth in detail below. Thus, preferred CBPs of the invention are antigenic, and more preferably immunogenic.

A molecule is "antigenic" when it is capable of specifically interacting with an antigen recognition molecule of the immune system, such as an immunoglobulin (antibody) or T cell antigen receptor. An antigenic polypeptide contains at least about 5, and preferably at least about 10, amino acids. An antigenic portion of a molecule can be that portion that is immunodominant for antibody or T cell receptor recognition, or it can be a portion used to generate an antibody to the molecule by conjugating the antigenic portion to a carrier molecule for immunization. A molecule that is antigenic need not be itself immunogenic, *i.e.*, capable of eliciting a humoral immune response without a carrier.

An "antibody" for purposes of this invention is any immunoglobulin, including antibodies and fragments thereof, that binds a specific epitope on a CBP. The term encompasses polyclonal, monoclonal, and chimeric antibodies, the last mentioned described in further detail in U.S. Patent Nos. 4,816,397 and 4,816,567.

An "antibody combining site" is that structural portion of an antibody molecule comprised of heavy and light chain variable and hypervariable regions that specifically binds antigen.

The phrase "antibody molecule" in its various grammatical forms as used herein contemplates both an intact immunoglobulin molecule and an immunologically active portion of an immunoglobulin molecule.

According to the invention, CBP(s) produced recombinantly, by chemical synthesis, or purified from the natural bacterial source, and fragments or other derivatives or analogs thereof, including fusion proteins, may be used as an immunogen to generate anti-CBP antibodies. Such antibodies include but are not limited to

polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library.

- Various procedures known in the art may be used for the production of polyclonal antibodies to CBPs or derivatives or analogs thereof (*see, e.g., Antibodies -- A Laboratory Manual*, Harlow and Lane, eds., Cold Spring Harbor Laboratory Press: Cold Spring Harbor, Ne York, 1988). For the production of antibody, various host animals can be immunized by injection with CBP(s), or a derivative (*e.g.,* fragment or fusion protein) thereof, including but not limited to rabbits, mice, rats, sheep, goats, etc. In one embodiment, the CBP or fragment thereof can be conjugated to an immunogenic carrier, *e.g.,* bovine serum albumin (BSA) or keyhole limpet hemocyanin (KLH). Various adjuvants may be used to increase the immunological response, depending on the host species.
- 15 For preparation of monoclonal antibodies directed toward the CBP, or fragment, analog, or derivative thereof, any technique that provides for the production of antibody molecules by continuous cell lines in culture may be used (*see, e.g., Antibodies -- A Laboratory Manual*, Harlow and Lane, eds., Cold Spring Harbor Laboratory Press: Cold Spring Harbor, Ne York, 1988). These include but are not limited to the hybridoma technique originally developed by Kohler and Milstein (1975, *Nature* 256:495-497), as well as the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals utilizing recent technology (PCT/US90/02545). According to the invention, human antibodies may be used and can be obtained by using human hybridomas (Cote et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:2026-2030) or by transforming human B cells with EBV virus *in vitro* (Cole et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, pp. 77-96). In fact,

according to the invention, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, J. Bacteriol. 159:870; Neuberger et al., 1984, Nature 312:604-608; Takeda et al., 1985, Nature 314:452-454) by splicing the genes from a mouse antibody molecule specific for a CBP together with genes from

5 a human antibody molecule of appropriate biological activity can be used; such antibodies are within the scope of this invention. Such human or humanized chimeric antibodies are preferred for use in therapy of human diseases or disorders (described *infra*), since the human or humanized antibodies are much less likely than xenogenic antibodies to induce an immune response, in particular an allergic

10 response, themselves.

According to the invention, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce CBP-specific single chain antibodies. An additional embodiment of the invention utilizes the techniques

15 described for the construction of Fab expression libraries (Huse et al., 1989, Science 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for a CBP, or its derivatives, or analogs.

Antibody fragments which contain the idiotype of the antibody molecule can be

20 generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, and the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent.

25

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, *e.g.*, radioimmunoassay, ELISA (enzyme-linked immunosorbant assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion

30 assays, *in situ* immunoassays (using colloidal gold, enzyme or radioisotope labels,

for example), western blots, precipitation reactions, agglutination assays (*e.g.*, gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention. For example, to select antibodies which recognize a specific epitope of a CBP, one may assay generated hybridomas for a product which binds to a CBP fragment containing such epitope. For selection of an antibody specific to a CBP from a particular strain of Gram positive bacteria, particularly pneumococcus, one can select on the basis of positive binding with CBP expressed by or isolated from cells of that strain of bacteria.

The foregoing antibodies can be used in methods known in the art relating to the localization and activity of the CBP, *e.g.*, for Western blotting, imaging CBP polypeptide *in situ*, measuring levels thereof in appropriate physiological samples, etc.

In a specific embodiment, antibodies that agonize or antagonize the activity of CBP can be generated. Such antibodies can be tested using the assays described *supra* for characterizing CBPs.

The term "adjuvant" refers to a compound or mixture that enhances the immune response to an antigen. An adjuvant can serve as a tissue depot that slowly releases the antigen and also as a lymphoid system activator that non-specifically enhances the immune response (Hood et al., *Immunology, Second Ed.*, 1984, Benjamin/Cummings: Menlo Park, California, p. 384). Often, a primary challenge with an antigen alone, in the absence of an adjuvant, will fail to elicit a humoral or

cellular immune response. Adjuvants include, but are not limited to, complete Freund's adjuvant, incomplete Freund's adjuvant, saponin, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (*bacille Calmette-Guerin*) and *Corynebacterium parvum*. Preferably, the adjuvant is pharmaceutically acceptable.

Vaccination and Passive Immune Therapy

- 10 Active immunity against Gram positive bacteria, particularly pneumococcus, can be induced by immunization (vaccination) with an immunogenic amount of an CBP, or an antigenic derivative or fragment thereof, and an adjuvant, wherein the CBP, or antigenic derivative or fragment thereof, is the antigenic component of the vaccine.
- 15 The CBP alone or conjugated to a capsule or capsules cannot cause bacterial infection, and the active immunity elicited by vaccination with the protein according to the present invention can result in both an immediate immune response and in immunological memory, and thus provide long-term protection against infection by the bacterium. The CBPs of the present invention, or antigenic fragments thereof,
- 20 can be prepared in an admixture with an adjuvant to prepare a vaccine. Preferably, the CBP, or derivative or fragment thereof, used as the antigenic component of the vaccine is an adhesin. More preferably, the CBP, or derivative or fragment thereof, used as the antigenic component of the vaccine is an antigen common to all or many strains of a species of Gram positive bacteria, or common to closely related
- 25 species of bacteria. Most preferably, the antigenic component of the vaccine is an adhesin that is a common antigen.

Selection of an adjuvant depends on the subject to be vaccinated. Preferably, a pharmaceutically acceptable adjuvant is used. For example, a vaccine for a human

30 should avoid oil or hydrocarbon emulsion adjuvants, including complete and

incomplete Freund's adjuvant. One example of an adjuvant suitable for use with humans is alum (alumina gel). A vaccine for an animal, however, may contain adjuvants not appropriate for use with humans.

- 5 An alternative to a traditional vaccine comprising an antigen and an adjuvant involves the direct *in vivo* introduction of DNA encoding the antigen into tissues of a subject for expression of the antigen by the cells of the subject's tissue. Such vaccines are termed herein "nucleic acid-based vaccines." Since the *CBP* gene by definition contains a signal sequence, expression of the gene in cells of the tissue
- 10 results in secretion of membrane association of the expressed protein. Alternatively, the expression vector can be engineered to contain an autologous signal sequence instead of the *CBP* signal sequence. For example, a naked DNA vector (*see, e.g.*, Ulmer et al., 1993, Science 259:1745-1749), a DNA vector transporter (*e.g.*, Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988,
- 15 J. Biol. Chem. 263:14621-14624; Hartmut et al., Canadian Patent Application No. 2,012,311, filed March 15, 1990), or a viral vector containing the desired *CBP* gene can be injected into tissue. Suitable viral vectors include retroviruses that are packaged in cells with amphotropic host range (*see* Miller, 1990, Human Gene Ther. 1:5-14; Ausubel et al., *Current Protocols in Molecular Biology*, § 9), and
- 20 attenuated or defective DNA virus, such as but not limited to herpes simplex virus (HSV) (*see, e.g.*, Kaplitt et al., 1991, Molec. Cell. Neurosci. 2:320-330), papillomavirus, Epstein Barr virus (EBV), adenovirus (*see, e.g.*, Stratford-Perricaudet et al., 1992, J. Clin. Invest. 90:626-630), adeno-associated virus (AAV) (*see, e.g.*, Samulski et al., 1987, J. Virol. 61:3096-3101; Samulski et al., 1989, J.
- 25 Virol. 63:3822-3828), and the like. Defective viruses, which entirely or almost entirely lack viral genes, are preferred. Defective virus is not infective after introduction into a cell.

Vectors containing the nucleic acid-based vaccine of the invention can be introduced

30 into the desired host by methods known in the art, *e.g.*, transfection,

electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, *e.g.*, Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624; Hartmut et al., Canadian Patent
 5 Application No. 2,012,311, filed March 15, 1990).

Either vaccine of the invention, *i.e.*, a vaccines comprising an CBP antigen or antigenic derivative or fragment thereof, or an *CBP* nucleic acid vaccine, can be administered via any parenteral route, including but not limited to intramuscular,
 10 intraperitoneal, intravenous, and the like. Preferably, since the desired result of vaccination is to elucidate an immune response to the antigen, and thereby to the pathogenic organism, administration directly, or by targeting or choice of a viral vector, indirectly, to lymphoid tissues, *e.g.*, lymph nodes or spleen. Since immune cells are continually replicating, they are ideal target for retroviral vector-based
 15 nucleic acid vaccines, since retroviruses require replicating cells.

Passive immunity can be conferred to an animal subject suspected of suffering an infection with a Gram positive bacterium, preferably streptococcal, and more preferably pneumoccal, by administering antiserum, polyclonal antibodies, or a
 20 neutralizing monoclonal antibody against a choline binding protein of the invention to the patient. Although passive immunity does not confer long term protection, it can be a valuable tool for the treatment of a bacterial infection of a subject who has not been vaccinated. Passive immunity is particularly important for the treatment of antibiotic resistant strains of Gram positive bacteria, since no other therapy may be
 25 available. Preferably, the antibodies administered for passive immune therapy are autologous antibodies. For example, if the subject is a human, preferably the antibodies are of human origin or have been "humanized," in order to minimize the possibility of an immune response against the antibodies. In a specific example, *infra*, passive immunity completely protects against lethal bacterial infection.

An analogous therapy to passive immunization is administration of an amount of an CBP protein adhesin sufficient to inhibit adhesion of the bacterium to its target cell. The required amount can be determined by one of ordinary skill using standard techniques.

5

The active or passive vaccines of the invention, or the administration of an adhesin, can be used to protect an animal subject from infection of a Gram positive bacteria, preferably streptococcus, and more preferably, pneumococcus. Thus, a vaccine of the invention can be used in birds, such as chickens, turkeys, and pets; in

10

mammals, preferably a human, although the vaccines of the invention are contemplated for use in other mammalian species, including but not limited to domesticated animals (canine and feline); farm animals (bovine, ovine, equine, caprine, porcine, and the like); rodents; and undomesticated animals.

15

Diagnosis of a Gram Positive Bacterial Infection

The antibodies of the present invention that can be generated against the CBPs from Gram positive bacteria are valuable reagents for the diagnosis of an infection with a Gram positive microorganism, particularly a pneumococcus. Presently, diagnosis of infection with a Gram positive bacterium is difficult. According to the invention,

20

the presence of Gram positive bacteria in a sample from a subject suspected of having an infection with a Gram positive bacterium can be detected by detecting binding of an antibody to an CBP to bacteria in or from the sample. In one aspect of the invention, the antibody can be specific for a unique strain or a limited number of strains of the bacterium, thus allowing for diagnosis of infection with

25

that particular strain (or strains). Alternatively, the antibody can be specific for many or all strains of a bacterium, thus allowing for diagnosis of infection with that species.

Diagnosis of infection with a Gram positive bacterium can use any immunoassay

30

format known in the art, as desired. Many possible immunoassay formats are

described in the section entitled "Antibodies to CBPs." The antibodies can be labeled for detection *in vitro*, *e.g.*, with labels such as enzymes, fluorophores, chromophores, radioisotopes, dyes, colloidal gold, latex particles, and chemiluminescent agents. Alternatively, the antibodies can be labeled for detection

5 *in vivo*, *e.g.*, with radioisotopes (preferably technetium or iodine); magnetic resonance shift reagents (such as gadolinium and manganese); or radio-opaque reagents.

In specific embodiments, the presence of CBP in or on cells can be ascertained by

10 the usual immunological procedures applicable to such determinations. A number of useful procedures are known. The procedures and their application are all familiar to those skilled in the art and accordingly may be utilized within the scope of the present invention. For example, a "competitive" procedure is described in U.S. Patent Nos. 3,654,090 and 3,850,752. A "sandwich" procedure, is described

15 in U.S. Patent Nos. RE 31,006 and 4,016,043. Still other procedures are known such as the "double antibody," or "DASP" procedure.

In each instance, the CBP forms complexes with one or more antibody(ies) or binding partners and one member of the complex is labeled with a detectable label.

20 The fact that a complex has formed and, if desired, the amount thereof, can be determined by known methods applicable to the detection of labels.

The labels most commonly employed for these studies are radioactive elements, enzymes, chemicals which fluoresce when exposed to ultraviolet light, and others.

25

A number of fluorescent materials are known and can be utilized as labels. These include, for example, fluorescein, rhodamine, auramine, Texas Red, AMCA blue and Lucifer Yellow. A particular detecting material is anti-rabbit antibody prepared in goats and conjugated with fluorescein through an isothiocyanate.

30

The CBP or its binding partner(s) can also be labeled with a radioactive element or with an enzyme. The radioactive label can be detected by any of the currently available counting procedures. The preferred isotope may be selected from ^3H , ^{14}C , ^{32}P , ^{35}S , ^{36}Cl , ^{51}Cr , ^{57}Co , ^{58}Co , ^{59}Fe , ^{90}Y , ^{125}I , ^{131}I , and ^{186}Re .

5

Enzyme labels are likewise useful, and can be detected by any of the presently utilized colorimetric, spectrophotometric, fluorospectrophotometric, amperometric or gasometric techniques. The enzyme is conjugated to the selected particle by reaction with bridging molecules such as carbodiimides, diisocyanates,

10 glutaraldehyde and the like. Many enzymes which can be used in these procedures are known and can be utilized. The preferred are peroxidase, β -glucuronidase, β -D-glucosidase, β -D-galactosidase, urease, glucose oxidase plus peroxidase and alkaline phosphatase. U.S. Patent Nos. 3,654,090; 3,850,752; and 4,016,043 are referred to by way of example for their disclosure of alternate labeling material and
15 methods.

In a further embodiment of this invention, commercial test kits suitable for use by a medical specialist may be prepared to determine the presence or absence of predetermined binding activity or predetermined binding activity capability to
20 suspected target cells. In accordance with the testing techniques discussed above, one class of such kits will contain at least the labeled CBP or its binding partner, for instance an antibody specific thereto, and directions, of course, depending upon the method selected, e.g., "competitive," "sandwich," "DASP" and the like. The kits may also contain peripheral reagents such as buffers, stabilizers, etc.

25

Accordingly, a test kit may be prepared for the demonstration of the presence or capability of cells for predetermined bacterial binding activity, comprising:

(a) a predetermined amount of at least one labeled immunochemically reactive component obtained by the direct or indirect attachment of the present CBP factor
30 or a specific binding partner thereto, to a detectable label;

- (b) other reagents; and
- (c) directions for use of said kit.

More specifically, the diagnostic test kit may comprise:

- 5 (a) a known amount of the CBP as described above (or a binding partner) generally bound to a solid phase to form an immunosorbent, or in the alternative, bound to a suitable tag, or plural such end products, etc. (or their binding partners) one of each;
- (b) if necessary, other reagents; and
- 10 (c) directions for use of said test kit.

In a further variation, the test kit may be prepared and used for the purposes stated above, which operates according to a predetermined protocol (e.g. "competitive," "sandwich," "double antibody," etc.), and comprises:

- 15 (a) a labeled component which has been obtained by coupling the CBP to a detectable label;
- (b) one or more additional immunochemical reagents of which at least one reagent is a ligand or an immobilized ligand, which ligand is selected from the group consisting of:
 - 20 (i) a ligand capable of binding with the labeled component (a);
 - (ii) a ligand capable of binding with a binding partner of the labeled component (a);
 - (iii) a ligand capable of binding with at least one of the component(s) to be determined; and
 - 25 (iv) a ligand capable of binding with at least one of the binding partners of at least one of the component(s) to be determined; and
- (c) directions for the performance of a protocol for the detection and/or determination of one or more components of an immunochemical reaction between the CBP and a specific binding partner thereto.

Alternatively, the nucleic acids and sequences thereof of the invention can be used in the diagnosis of infection with a Gram positive bacterium. For example, the *CBP* genes or hybridizable fragments thereof can be used for *in situ* hybridization with a sample from a subject suspected of harboring an infection of Gram positive bacteria. In another embodiment, specific gene segments of a Gram positive bacterium can be identified using PCR amplification with probes based on the *CBP* genes of the invention. In one aspect of the invention, the hybridization with a probe or with the PCR primers can be performed under stringent conditions, or with a sequence specific for a unique strain or a limited number of strains of the bacterium, or both, thus allowing for diagnosis of infection with that particular strain (or strains). Alternatively, the hybridization can be under less stringent conditions, or the sequence may be homologous in any or all strains of a bacterium, thus allowing for diagnosis of infection with that species.

15 Therapeutic Compositions and Vaccines Comprising CBPs

As noted above, the present invention provides therapeutic compositions comprising antibodies to CBPs (*i.e.*, passive immune therapy), anti-CBP vaccines (whether comprised of CBPs in an adjuvant, or a nucleic acid vaccine), CBPs to compete with bacterial CBPs for pathogenic activities, such as adherence to host cells, peptides, such as Fn5, or antibodies to Fn5. Preferably, any such composition is a pharmaceutical composition comprising the active component (antibody, vaccine, or CBP) in a pharmaceutically acceptable carrier.

The preparation of therapeutic compositions which contain an active component is well understood in the art. Typically, such compositions are prepared as injectables, either as liquid solutions or suspensions, however, solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified. The active therapeutic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose,

glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents which enhance the effectiveness of the active ingredient.

5

A active component can be formulated into the therapeutic composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

10
15

A composition comprising "A" (where "A" is a single protein, DNA molecule, vector, etc.) is substantially free of "B" (where "B" comprises one or more contaminating proteins, DNA molecules, vectors, etc.) when at least about 75% by weight of the proteins, DNA, vectors (depending on the category of species to which A and B belong) in the composition is "A". Preferably, "A" comprises at least about 90% by weight of the A+B species in the composition, most preferably at least about 99% by weight.

20

The phrase "pharmaceutically acceptable" refers to molecular entities and compositions that are physiologically tolerable and do not typically produce an allergic or similar untoward reaction, such as gastric upset, dizziness and the like, when administered to a human. Preferably, as used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally

25
30

recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water or aqueous solution saline solutions and aqueous dextrose and glycerol solutions are preferably employed as carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin.

10

The phrase "therapeutically effective amount" is used herein to mean an amount sufficient to reduce by at least about 15 percent, preferably by at least 50 percent, more preferably by at least 90 percent, and most preferably prevent, a clinically significant deficit in the activity, function and response of the host. Alternatively, a therapeutically effective amount is sufficient to cause an improvement in a clinically significant condition in the host. In the context of the present invention, a deficit in the response of the host is evidenced by continuing or spreading bacterial infection. An improvement in a clinically significant condition in the host includes a decrease in bacterial load, clearance of bacteria from colonized host cells, reduction in fever or inflammation associated with infection, or a reduction in any symptom associated with the bacterial infection.

15
20

According to the invention, the component or components of a therapeutic composition of the invention may be introduced parenterally, transmucosally, *e.g.*, orally, nasally, pulmonarily, or rectally, or transdermally. Preferably, administration is parenteral, *e.g.*, via intravenous injection, and also including, but is not limited to, intra-arteriole, intramuscular, intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial administration. Oral or pulmonary delivery may be preferred to activate mucosal immunity; since pneumococci generally colonize the nasopharyngeal and pulmonary mucosa, mucosal immunity

25
30

may be a particularly effective preventive treatment. The term "unit dose" when used in reference to a therapeutic composition of the present invention refers to physically discrete units suitable as unitary dosage for humans, each unit containing a predetermined quantity of active material calculated to produce the desired
 5 therapeutic effect in association with the required diluent; *i.e.*, carrier, or vehicle.

In another embodiment, the active compound can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and
 10 Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

In yet another embodiment, the therapeutic compound can be delivered in a controlled release system. For example, the polypeptide may be administered using
 15 intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of*
 20 *Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J. *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann. Neurol.* 25:351 (1989); Howard et al.,
 25 *J. Neurosurg.* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, *i.e.*, the brain, thus requiring only a fraction of the systemic dose (see, *e.g.*, Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)). Preferably, a controlled release device is introduced into a subject in proximity of the site of
 30 inappropriate immune activation or a tumor.

Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

- A subject in whom administration of an active component as set forth above is an effective therapeutic regimen for a bacterial infection is preferably a human, but can be any animal. Thus, as can be readily appreciated by one of ordinary skill in the art, the methods and pharmaceutical compositions of the present invention are particularly suited to administration to any animal, particularly a mammal, and including, but by no means limited to, domestic animals, such as feline or canine subjects, farm animals, such as but not limited to bovine, equine, caprine, ovine, and porcine subjects, wild animals (whether in the wild or in a zoological garden), research animals, such as mice, rats, rabbits, goats, sheep, pigs, dogs, cats, etc., *i.e.*, for veterinary medical use.
- 15 In the therapeutic methods and compositions of the invention, a therapeutically effective dosage of the active component is provided. A therapeutically effective dosage can be determined by the ordinary skilled medical worker based on patient characteristics (age, weight, sex, condition, complications, other diseases, etc.), as is well known in the art. Furthermore, as further routine studies are conducted,
- 20 more specific information will emerge regarding appropriate dosage levels for treatment of various conditions in various patients, and the ordinary skilled worker, considering the therapeutic context, age and general health of the recipient, is able to ascertain proper dosing. Generally, for intravenous injection or infusion, dosage may be lower than for intraperitoneal, intramuscular, or other route of
- 25 administration. The dosing schedule may vary, depending on the circulation half-life, and the formulation used. The compositions are administered in a manner compatible with the dosage formulation in the therapeutically effective amount. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual. However, suitable
- 30 dosages may range from about 0.1 to 20, preferably about 0.5 to about 10, and

more preferably one to several, milligrams of active ingredient per kilogram body weight of individual per day and depend on the route of administration. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration. Alternatively, continuous intravenous infusion sufficient to maintain concentrations of ten nanomolar to ten micromolar in the blood are contemplated.

Administration with other compounds. For treatment of a bacterial infection, one may administer the present active component in conjunction with one or more pharmaceutical compositions used for treating bacterial infection, including but not limited to (1) antibiotics; (2) soluble carbohydrate inhibitors of bacterial adhesion; (3) other small molecule inhibitors of bacterial adhesion; (4) inhibitors of bacterial metabolism, transport, or transformation; (5) stimulators of bacterial lysis, or (6) anti-bacterial antibodies or vaccines directed at other bacterial antigens. Other potential active components include anti-inflammatory agents, such as steroids and non-steroidal anti-inflammatory drugs. Administration may be simultaneous (for example, administration of a mixture of the present active component and an antibiotic), or may be *in seriatim*.

Accordingly, in specific embodiment, the therapeutic compositions may further include an effective amount of the active component, and one or more of the following active ingredients: an antibiotic, a steroid, etc. Exemplary formulations are given below:

Formulations

Intravenous Formulation I

<u>Ingredient</u>	<u>mg/ml</u>
cefotaxime	250.0

	CBP	10.0
	dextrose USP	45.0
	sodium bisulfite USP	3.2
	edetate disodium USP	0.1
5	water for injection q.s.a.d.	1.0 ml

Intravenous Formulation II

	<u>Ingredient</u>	<u>mg/ml</u>
	ampicillin	250.0
10	CBP	10.0
	sodium bisulfite USP	3.2
	disodium edetate USP	0.1
	water for injection q.s.a.d.	1.0 ml

15 Intravenous Formulation III

	<u>Ingredient</u>	<u>mg/ml</u>
	gentamicin (charged as sulfate)	40.0
	CBP	10.0
	sodium bisulfite USP	3.2
20	disodium edetate USP	0.1
	water for injection q.s.a.d.	1.0 ml

Intravenous Formulation IV

	<u>Ingredient</u>	<u>mg/ml</u>
25	CBP	10.0
	dextrose USP	45.0
	sodium bisulfite USP	3.2
	edetate disodium USP	0.1
	water for injection q.s.a.d.	1.0 ml

Intravenous Formulation V

<u>Ingredient</u>	<u>mg/ml</u>
CBP antagonist	5.0
sodium bisulfite USP	3.2
5 disodium edetate USP	0.1
water for injection q.s.a.d.	1.0 ml

As used herein, "pg" means picogram, "ng" means nanogram, "ug" or " μ g" mean microgram, "mg" means milligram, "ul" or " μ l" mean microliter, "ml" means
 10 milliliter, "l" means liter.

Thus, in a specific instance where it is desired to reduce or inhibit the infection resulting from CBP-mediated binding of bacteria to a host cell, CBP or an antibody thereto, or a ligand thereof or an antibody to that ligand, such as Fn5, could be
 15 introduced to block the interaction of CBP present on bacteria with the host cell.

As discussed earlier, the CBPs or antibodies thereto may be prepared in pharmaceutical compositions, with a suitable carrier and at a strength effective for administration by various means to a patient experiencing an adverse medical
 20 condition associated with specific bacterial infection for the treatment thereof. A variety of administrative techniques may be utilized, among them parenteral techniques such as subcutaneous, intravenous and intraperitoneal injections, catheterizations and the like. Average quantities of the CBPs or their subunits may vary and in particular should be based upon the recommendations and prescription
 25 of a qualified physician or veterinarian.

Pulmonary Delivery

Also contemplated herein is pulmonary delivery of the present adhesion inhibitory agent (or derivatives thereof), of the invention, selected from the group consisting
 30 of a choline binding protein, an antibody to a choline binding protein, an enolase,

hindered cationic small molecules (such as lysine, choline, arginine, etc.), the peptide WQPPRARI (SEQ ID NO:11), and antibody specific for an epitope having the amino acid sequence WQPPRARI (SEQ ID NO:11). The adhesion inhibitory agent (or derivative) is delivered to the lungs of a mammal, where it can interfere

5 with bacterial, *i.e.*, streptococcal, and preferably pneumococcal binding to host cells. In a specific embodiment, such an adhesion inhibitory agent inhibits binding of the streptococcal enolase to fibronectin. Other reports of preparation of proteins for pulmonary delivery are found in the art [Adjei et al. *Pharmaceutical Research*, 7:565-569 (1990); Adjei et al., *International Journal of Pharmaceutics*, 63:135-144

10 (1990) (leuprolide acetate); Braquet et al., *Journal of Cardiovascular Pharmacology*, 13(suppl. 5):143-146 (1989) (endothelin-1); Hubbard et al., *Annals of Internal Medicine*, Vol. III, pp. 206-212 (1989) (α 1-antitrypsin); Smith et al., *J. Clin. Invest.* 84:1145-1146 (1989) (α -1-proteinase); Oswein et al., "Aerosolization of Proteins", *Proceedings of Symposium on Respiratory Drug*

15 *Delivery II*, Keystone, Colorado, March, (1990) (recombinant human growth hormone); Debs et al., *J. Immunol.* 140:3482-3488 (1988) (interferon- γ and tumor necrosis factor alpha); Platz et al., U.S. Patent No. 5,284,656 (granulocyte colony stimulating factor)]. A method and composition for pulmonary delivery of drugs is described in U.S. Patent No. 5,451,569, issued September 19, 1995 to Wong et al.

20 Contemplated for use in the practice of this invention are a wide range of mechanical devices designed for pulmonary delivery of therapeutic products, including but not limited to nebulizers, metered dose inhalers, and powder inhalers, all of which are familiar to those skilled in the art. With regard to construction of

25 the delivery device, any form of aerosolization known in the art, including but not limited to spray bottles, nebulization, atomization or pump aerosolization of a liquid formulation, and aerosolization of a dry powder formulation, can be used in the practice of the invention.

Some specific examples of commercially available devices suitable for the practice of this invention are the Ultravent nebulizer, manufactured by Mallinckrodt, Inc., St. Louis, Missouri; the Acorn II nebulizer, manufactured by Marquest Medical Products, Englewood, Colorado; the Ventolin metered dose inhaler, manufactured by Glaxo Inc., Research Triangle Park, North Carolina; and the Spinhaler powder inhaler, manufactured by Fisons Corp., Bedford, Massachusetts.

All such devices require the use of formulations suitable for the dispensing of adhesion inhibitory agent (or derivative). Typically, each formulation is specific to the type of device employed and may involve the use of an appropriate propellant material, in addition to the usual diluents, adjuvants and/or carriers useful in therapy. Also, the use of liposomes, microcapsules or microspheres, inclusion complexes, or other types of carriers is contemplated. Chemically modified adhesion inhibitory agent may also be prepared in different formulations depending on the type of chemical modification or the type of device employed.

Formulations suitable for use with a nebulizer, either jet or ultrasonic, will typically comprise adhesion inhibitory agent (or derivative) dissolved in water at a concentration of about 0.1 to 25 mg of biologically active adhesion inhibitory agent per ml of solution. The formulation may also include a buffer and a simple sugar (e.g., for adhesion inhibitory agent stabilization and regulation of osmotic pressure). The nebulizer formulation may also contain a surfactant, to reduce or prevent surface induced aggregation of the adhesion inhibitory agent caused by atomization of the solution in forming the aerosol.

25

Formulations for use with a metered-dose inhaler device will generally comprise a finely divided powder containing the adhesion inhibitory agent (or derivative) suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as a chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including

trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol, and 1,1,1,2-tetrafluoroethane, or combinations thereof. Suitable surfactants include sorbitan trioleate and soya lecithin. Oleic acid may also be useful as a surfactant.

- 5 The liquid aerosol formulations contain adhesion inhibitory agent and a dispersing agent in a physiologically acceptable diluent. The dry powder aerosol formulations of the present invention consist of a finely divided solid form of adhesion inhibitory agent and a dispersing agent. With either the liquid or dry powder aerosol formulation, the formulation must be aerosolized. That is, it must be broken down
10 into liquid or solid particles in order to ensure that the aerosolized dose actually reaches the mucous membranes of the nasal passages or the lung. The term "aerosol particle" is used herein to describe the liquid or solid particle suitable for nasal or pulmonary administration, *i.e.*, that will reach the mucous membranes. Other considerations, such as construction of the delivery device, additional
15 components in the formulation, and particle characteristics are important. These aspects of pulmonary administration of a drug are well known in the art, and manipulation of formulations, aerosolization means and construction of a delivery device require at most routine experimentation by one of ordinary skill in the art.
- 20 In a particular embodiment, the mass median dynamic diameter will be 5 micrometers or less in order to ensure that the drug particles reach the lung alveoli [Wearley, L.L., *Crit. Rev. in Ther. Drug Carrier Systems* 8:333 (1991)].

- Systems of aerosol delivery, such as the pressurized metered dose inhaler and the
25 dry powder inhaler are disclosed in Newman, S.P., *Aerosols and the Lung*, Clarke, S.W. and Davia, D. editors, pp. 197-22 and can be used in connection with the present invention.

- In a further embodiment, as discussed in detail *infra*, an aerosol formulation of the
30 present invention can include other therapeutically or pharmacologically active

ingredients in addition to adhesion inhibitory agent, such as but not limited to an antibiotic, a steroid, a non-steroidal anti-inflammatory drug, etc.

Liquid Aerosol Formulations. The present invention provides aerosol formulations
5 and dosage forms for use in treating subjects suffering from bacterial, *e.g.*,
streptococcal, in particularly pneumococcal, infection. In general such dosage
forms contain adhesion inhibitory agent in a pharmaceutically acceptable diluent.
Pharmaceutically acceptable diluents include but are not limited to sterile water,
saline, buffered saline, dextrose solution, and the like. In a specific embodiment, a
10 diluent that may be used in the present invention or the pharmaceutical formulation
of the present invention is phosphate buffered saline, or a buffered saline solution
generally between the pH 7.0-8.0 range, or water.

The liquid aerosol formulation of the present invention may include, as optional
15 ingredients, pharmaceutically acceptable carriers, diluents, solubilizing or
emulsifying agents, surfactants and excipients.

The formulation may include a carrier. The carrier is a macromolecule which is
soluble in the circulatory system and which is physiologically acceptable where
20 physiological acceptance means that those of skill in the art would accept injection
of said carrier into a patient as part of a therapeutic regime. The carrier preferably
is relatively stable in the circulatory system with an acceptable plasma half life for
clearance. Such macromolecules include but are not limited to Soya lecithin, oleic
acid and sorbitan trioleate, with sorbitan trioleate preferred.

25

The formulations of the present embodiment may also include other agents useful
for pH maintenance, solution stabilization, or for the regulation of osmotic
pressure. Examples of the agents include but are not limited to salts, such as
sodium chloride, or potassium chloride, and carbohydrates, such as glucose,
30 galactose or mannose, and the like.

The present invention further contemplates liquid aerosol formulations comprising adhesion inhibitory agent and another therapeutically effective drug, such as an antibiotic, a steroid, a non-steroidal anti-inflammatory drug, etc.

- 5 *Aerosol Dry Powder Formulations.* It is also contemplated that the present aerosol formulation can be prepared as a dry powder formulation comprising a finely divided powder form of adhesion inhibitory agent and a dispersant.

Formulations for dispensing from a powder inhaler device will comprise a finely
 10 divided dry powder containing adhesion inhibitory agent (or derivative) and may also include a bulking agent, such as lactose, sorbitol, sucrose, or mannitol in amounts which facilitate dispersal of the powder from the device, *e.g.*, 50 to 90% by weight of the formulation. The adhesion inhibitory agent (or derivative) should most advantageously be prepared in particulate form with an average particle size of
 15 less than 10 mm (or microns), most preferably 0.5 to 5 mm, for most effective delivery to the distal lung.

In another embodiment, the dry powder formulation can comprise a finely divided dry powder containing adhesion inhibitory agent, a dispersing agent and also a
 20 bulking agent. Bulking agents useful in conjunction with the present formulation include such agents as lactose, sorbitol, sucrose, or mannitol, in amounts that facilitate the dispersal of the powder from the device.

The present invention further contemplates dry powder formulations comprising
 25 adhesion inhibitory agent and another therapeutically effective drug, such as an antibiotic, a steroid, a non-steroidal anti-inflammatory drug, etc.

Identification of Small Molecule Antagonists of CBPs

Identification and isolation of a gene encoding a choline binding protein of the
 30 invention provides for expression of the protein in quantities greater than can be

isolated from natural sources, or in indicator cells that are specially engineered to indicate the activity of a CBP expressed after transfection or transformation of the cells. According, in addition to rational design of agonists and antagonists based on the structure of a choline binding protein, the present invention contemplates an
5 alternative method for identifying specific ligands of a choline binding protein using various screening assays known in the art. Examples of small molecule antagonists of CBPs include lysine, choline, the pentapeptide having SEQ ID NO:11; it is likely that analogs of these types of molecules, such as arginine, will also inhibit bacterial adhesion to host tissue, particularly fibronectin. Thus, the present invention
10 generally provides a small molecule CBP binding antagonist, which is a hindered cationic molecule, preferably a hindered amine.

Any screening technique known in the art can be used to screen for choline binding protein agonists or antagonists. The present invention contemplates screens for
15 small molecule ligands or ligand analogs and mimics, as well as screens for natural ligands that bind to and agonize or antagonize activities of choline binding protein *in vivo*, particularly adhesion of bacteria to host cells or tissues mediated by the choline binding protein.

20 Knowledge of the primary sequence of the CBP, and the similarity of that sequence with proteins of known function, can provide an initial clue as the inhibitors or antagonists of the protein. Identification and screening of antagonists is further facilitated by determining structural features of the protein, *e.g.*, using X-ray crystallography, neutron diffraction, nuclear magnetic resonance spectrometry, and
25 other techniques for structure determination. These techniques provide for the rational design or identification of agonists and antagonists.

Another approach uses recombinant bacteriophage to produce large libraries. Using the "phage method" (Scott and Smith, 1990, Science 249:386-390; Cwirla, et al.,
30 1990, Proc. Natl. Acad. Sci., 87:6378-6382; Devlin et al., 1990, Science, 249:404-

406), very large libraries can be constructed (10^6 - 10^8 chemical entities). A second approach uses primarily chemical methods, of which the Geysen method (Geysen et al., 1986, Molecular Immunology 23:709-715; Geysen et al. 1987, J. Immunologic Method 102:259-274) and the recent method of Fodor et al. (1991, Science 251, 5 767-773) are examples. Furka et al. (1988, 14th International Congress of Biochemistry, Volume 5, Abstract FR:013; Furka, 1991, Int. J. Peptide Protein Res. 37:487-493), Houghton (U.S. Patent No. 4,631,211, issued December 1986) and Rutter et al. (U.S. Patent No. 5,010,175, issued April 23, 1991) describe methods to produce a mixture of peptides that can be tested as agonists or 10 antagonists.

In another aspect, synthetic libraries (Needels et al., 1993, "Generation and screening of an oligonucleotide encoded synthetic peptide library," Proc. Natl. Acad. Sci. USA 90:10700-4; Ohlmeyer et al., 1993, Proc. Natl. Acad. Sci. USA 15 90:10922-10926; Lam et al., International Patent Publication No. WO 92/00252; Kocis et al., International Patent Publication No. WO 9428028, each of which is incorporated herein by reference in its entirety), and the like can be used to screen for choline binding protein ligands according to the present invention.

20 The screening can be performed with recombinant cells that express the choline binding protein, or alternatively, using purified protein, *e.g.*, produced recombinantly, as described above. For example, the ability of labeled, soluble or solubilized choline binding protein that includes the ligand-binding portion of the molecule, to bind ligand can be used to screen libraries, as described in the 25 foregoing references.

The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

5

EXAMPLES: PRELIMINARY CONSIDERATIONS

The present invention identifies a family of CBPs which are surface exposed proteins and serve as adhesins critical for the binding of pneumococcus to eukaryotic cells found at the sites of infection. The invention also identifies and
 10 characterizes a novel CBP, CbpA, as an adhesin and a determinant of virulence. Furthermore, the present invention demonstrates that these proteins are immunogenic and thus may also function as protective antigens.

Using choline bound to agarose beads, a family of at least 12 bacterial choline
 15 binding proteins (CBPs), including LytA, was purified from a serotype II, *PspA* strain of pneumococcus. Labeling of whole bacteria with fluorescein isothiocyanate (FITC) suggested that four of these CBPs were surface-exposed while human convalescent sera cross-reacted with five of the CBPs. It was previously shown that pneumococcus binds to type II lung cells (LC) and endothelial cells (EC) of the
 20 peripheral vasculature. Compared to controls, the CBP fraction blocked pneumococcal adherence by 45% to LC and 89% to EC in a dose-dependent manner. The CBP fraction was used to immunize rabbits, and polyclonal antibodies were used for passive immunization studies. N-terminal and internal amino acid sequence obtained from three of the CBPs showed no similarity to any known
 25 protein sequences. Moreover, the predominant member of this family of proteins, CbpA, is a 112 kDa surface exposed protein that reacted with human convalescent sera. Sequence analysis of the corresponding gene showed a unique N-terminal sequence and six C-terminal choline binding domains in the C-terminal region.

EXAMPLE 1: IDENTIFICATION OF PNEUMOCOCCAL
CHOLINE BINDING PROTEINS

Materials and Methods

- 5 *Strains.* Bacterial strains used included (1) R6, an unencapsulated bacterial strain derived from D39 (type 2), unencapsulated, (2) AII encapsulated, (3) LM91 derived from D39 (encapsulated), which is PspA⁻, and (4) Lyt⁻ derived from D39, unencapsulated.
- 10 *Cell culture and choline binding protein purification.* Cultures of 0.1 ml bacteria LM91 (-80°C stock) were cultivated in 10 ml C + Y semisynthetic medium for 5 hours at 37°C. This 10 ml culture was then used to inoculate 400 ml C + Y, and this culture was incubated for 5 hours at 37°C in 5% CO₂ to an OD₆₂₀ of 0.6. To this mixture was added 4 ml choline agarose beads (CAB), and this was incubated
- 15 30 minutes at 37°C in 5% CO₂, with occasional shaking. The bacteria and beads were centrifuged at 4°C for 15 minutes at 8000 x g, and the pellet was resuspended in 50 ml rest medium (C + Y). To this 50 ml was added 20 µg/ml Leupeptin, 100 µg/ml PMSF and 250 µl Triton X100 to a final concentration of 0.5%. This mixture was rotated for 20-30 minutes until the solution became clear, indicating
- 20 that all bacteria were lysed. The CAB were then washed on a glass filter with DPBS/1 M NaCl. The CAB were then incubated with 2 changes of 50 ml DPBS/1 M NaCl, and the beads were again washed on a glass filter.

- Purification of choline binding proteins.* Choline binding proteins (CBPs) were
- 25 eluted with 3 washes of 5 ml DPBS/10% choline. The eluted CBPs were concentrated on an Ultrafree-20 concentrator (Millipore) to 1.5 ml, and then dialyzed against PBS.

- Preparation of rabbit anti-CBP antiserum.* Choline binding proteins isolated from
- 30 the choline affinity column as described above were concentrated and used to immunize rabbits in a vaccine to generate an anti-CBP antiserum. New Zealand

White rabbits were immunized by intradermal injection in the back with 500 μ g purified CBPs in 1 ml of a 1:1 emulsion of buffer and Complete Freund's Adjuvant. The rabbits were boosted with a subcutaneous dorsal injection of 250 μ g of purified CBPs in 1 ml of 1:1 buffer to Incomplete Freund's Adjuvant at weeks 3, 6, 9, 12, and 15. Rabbits were pre-bled, and test bled approximately 10 days prior to each boost. Serum obtained at week 16 were used in all experiments.

The CBPs were examined by SDS-PAGE and Western blot using the following antibodies: (1) polyclonal anti-pneumococcal serum (rabbit or "ROB"); (2) anti-PspA monoclonal antibody; (3) anti-FITC monoclonal antibody; (4) convalescent human anti-pneumococcal sera and (5) polyclonal anti-CBP antibody. 9 CBPs were identified on SDS-PAGE/Western blot. Results are shown in Figures 1 and 2.

15 EXAMPLE 2: CHARACTERIZATION OF PNEUMOCOCCAL CHOLINE BINDING PROTEINS

The CBPs isolated by SDS-PAGE were subjected to N-terminal amino acid sequence analysis. The protein fraction obtained from a choline agarose column eluted from 0.5 M choline chloride was applied to an SDS polyacrylamide gel transferred to a PVDF membrane, and stained for protein. Individual bands were excised and N-terminal amino acid sequence obtained by Edman degradation. Internal sequence data was obtained from protease generated peptide fragments. Amino acid sequence data for 8 of the 9 CBPs were obtained as follows:

- 25 CBP112 XNEGSTQAATSSNMAKTEHRKAAKQVVDE (SEQ ID NO:1)
- CBP90 AREFSLEKTR (SEQ ID NO:2)
- CBP84 XREFSLEKTRNIGIMAHVDAGKT (SEQ ID NO:3)
- CBP80 XKXXWQXKQYLKEDGSQAANEXVFDTA (SEQ ID NO:4)
- CBP78 QKIIGIDLGTNSAVAVLEGTESKIIANPE (SEQ ID NO:5)
- 30 CBP70 XXXEVAKXSQDTTAS (SEQ ID NO:6)
- CBP60 XNERVKIVATLGPAVEGRG (SEQ ID NO:7)

CBP50 XIIXXVYAREVLDSRGNP (SEQ ID NO:8)

CBP112-Int1 EDRRNYHPTNTYK (SEQ ID NO:9)

CBP112-Int2 XDDQQAEDYA (SEQ ID NO:10)

5

None of the sequences had been previously identified, except that CBP84 was 85% identical to Elongation Factor G (EF-G) and CBP50 was similar to enolase.

10

EXAMPLE 3: PNEUMOCOCCAL CHOLINE BINDING
PROTEINS BLOCK ADHERENCE

The CBP fraction isolated in Example 1 was used in an adherence assays to determine its effect on pneumococcal adherence to type II lung cells (LC) and endothelial cells of the vasculature (EC) as described in International Patent No.

15 PCT/US95/07209, filed June 6, 1995 by Tuomanen and Cundell, which is specifically incorporated herein by reference in its entirety. The CBP fraction blocked pneumococcal adherence by 45% to LC and 89% to EC in a dose-dependent manner (Figures 3 and 4, respectively).

20

EXAMPLE 4: THE 50KD CHOLINE BINDING PROTEIN
MEDIATES PNEUMOCOCCAL ADHESION

Adherence to extracellular matrix proteins, such as fibronectin (see Figure 5), affords pathogens with a means to invade injured epithelia. *S. pneumoniae* is

25 known to adhere to immobilized fibronectin at a site within the C-terminal heparin binding domain [van der Flier, *et. al. Infect. Immun.* 63:4317-4322(1995)]. Others have reported that this region also binds the leukocyte integrin adhesion receptor $\alpha 4\beta 1$. It has also been demonstrated that vascular cell adhesion molecule-1 (VCAM-1) contains sequences homologous to IIICS that are the active binding sites
30 for $\alpha 4\beta 1$. This Example evaluates, *inter alia*, if *S. pneumoniae*, like $\alpha 4\beta 1$, cross-recognized fibronectin and VCAM-1.

Adherence of *S. pneumoniae* to immobilized fibronectin was inhibited 96% by preincubation of the bacteria with recombinant soluble VCAM-1. *S. pneumoniae* also adhered directly to VCAM-1 coated wells at a density of 75 ± 18 bacteria/0.25 mm². This represents 6% of the adherence of pneumococci to whole fibronectin.

- 5 *S. pneumoniae* adherence specificity was further characterized by evaluating adherence to fragments of the 33 kDa heparin binding domain of fibronectin. Various fragments of the 33 kDa domain were expressed as GST-fusion proteins. The results are shown in Figure 6. Bacteria adhered about equally to recombinant fragments 33/66, 51, and 15. Further studies using four synthetic peptides based on the III14 region identified the peptide FN5 (WQPPRARI (SEQ ID NO:11)) as able to support adherence of *S. pneumoniae* (Figure 7). Antibody to FN5 inhibited adherence of pneumococci to both whole fibronectin and VCAM-1 (92 and 69% respectively). This Example shows that *S. pneumoniae* binds by similar mechanisms to VCAM-1 and fibronectin, which may provide pneumococci with access to leukocyte trafficking pathways, promoting the progression of disease.
- 15

- One adhesin on the surface of *S. pneumoniae* is the 50 kDa choline binding protein. Preincubation of bacteria in a 10% choline solution, to displace surface proteins bound to choline, resulted in a greater than 50% decrease in binding of pneumococci to fibronectin (Figure 8). Bacteria grown in the presence of ethanolamine instead of choline, so that they do not display choline binding proteins on their surfaces, failed to bind to fibronectin (> 85% inhibition)(data not shown). Pneumococcal binding was decreased by greater than 95% by 1:10 to 1:5000 dilutions of the anti-choline antibody, TEPC15 (Figure 9). Preparations of choline binding proteins from *S. pneumoniae* competitively inhibited the adherence of pneumococci to fibronectin (75-90% inhibition)(data not shown).
- 20
 - 25

- To confirm the identity of the protein adhesin on the surface of the pneumococcus, whole cell French Press lysates and preparations of choline binding proteins were separated by SDS-PAGE and blotted to Immobilon PVDF membranes. The
- 30

membranes were then incubated with a fibronectin solution for several hours. After washing away unbound protein, bound fibronectin was detected by chemiluminescence. A band of approximately 50 kDa was observed in both the whole cell lysate and choline binding protein preparations. Amino-terminal protein sequencing of this band identified this protein as a homolog of the glycolytic enzyme, enolase (2-phospho-D-glycerate hydrolyase). In the 18 amino acid stretch analyzed by the BLAST algorithm, a 16 residue region was shown to have 93 % identity (15/16) with the enolase from *Bacillus subtilis* (Figure 10).

- 10 Addition of the glycolytic enzyme enolase (Sigma) to fibronectin coated wells prior to addition of bacteria resulted in inhibition of pneumococcal adherence to fibronectin (Figure 11). Inhibition was dose dependent. Maximum inhibition (98%) was observed with 1000 U/ml enolase. No direct adherence of pneumococcus to enolase coated wells was observed.

- 15 Enolase has been shown to adhere to plasminogen via its carboxy-terminal lysine residue. Adherence of *S. pneumoniae* to whole fibronectin is L-lysine dependent. Lysine was added to fibronectin coated wells 15 min prior to addition of bacteria. Pneumococcal adherence was inhibited 67%, 95%, and 99% by .1M, .5M, and 1M L-lysine, respectively (Figure 12). The lysine analog, 6-amino hexanoic acid, was not able to inhibit adherence.

- 25 *S. pneumoniae* French Press lysates were run over a column of fibronectin coated sepharose beads. Adherent proteins were competitively eluted with L-lysine (.01-.5 M) and analyzed by SDS-PAGE. A band of approximately 50 kDa was observed in all fractions (Figure 13). This band comigrated with commercial enolase (Sigma).

- 30 Partial sequence for the enolase gene and the deduced amino acid sequence been obtained (SEQ ID NOS: 10 and 20). Comparison of the DNA and deduced amino

acid sequences of pneumococcus and *B. subtilis* are shown in Figure 14 and 15, respectively. There is 74% identity at the DNA level, and 72% identity, 85% similarity, at the amino acid level. More complete sequence data for the CBP-50 enolase, lacking only a putative 42 base 5' stretch/14 amino acid N-terminal stretch
5 are shown in SEQ ID NOS:18 and 19, respectively.

In addition to binding to fibronectin and to VCAM, the enolase homolog has two other properties related to virulence. It binds to alpha 1 acid glycoprotein (AGP) which is decorated with carbohydrates that are the key determinants of binding to
10 activated eukaryotic cells. AGP was affixed to a column and a pneumococcal lysate was passed over the column and the proteins retained on the column were eluted. A 50 kDa protein eluted specifically and upon amino acid N-terminal sequencing was shown to be the enolase homolog. A second property of the enolase homolog is that it appears to be absent from avirulent pneumococci which have phase varied to
15 the opaque colonial morphology. When proteins eluted from pneumococci treated with 2% choline are compared between opaque and transparent strains, the opaque are missing a 112 kDa protein and the 50kD protein. This indicates that lack of virulence is associated with lack of expression of the enolase homolog and therefore enolase homolog is a true virulence determinant.

20

Materials and Methods

Bacterial strains and growth conditions. *S. pneumoniae* Type 2 (D39) and the isogenic, unencapsulated derivatives R6x and R6 were used. Pneumococcal strain
25 LM34 deleted for production of pneumococcal surface protein A (*pspA*-) has been described [McDaniel *et al.*, *Infect. Immun.*, **59**:222-9 (1991)]. Bacteria were grown on tryptic soy agar containing 5% sheep blood for 16-18 hours at 37°C in 5% CO₂ or in a candle extinction jar.

Fibronectin and derived proteolytic fragments. Fibronectin from human plasma was purchased from Sigma (St. Louis, MO) and from Gibco BRL (Grand Island, NY).

Fragments designated by molecular size are depicted schematically in Figure 5.

Proteolytic fragments purchased from Gibco BRL (indicated in parentheses in

- 5 Figure 5) included a tryptic 30kD collagen binding fragment, an α -chymotryptic 120kD cell binding fragment and an α -chymotryptic 40kD heparin binding fragment. Proteolytic fragments prepared following the method of Vercelotti et al [Vercelotti *et al.*, *J. Lab. Clin. Med.*, **103**:34-43 (1984); McCarthy *et al.*, 1986, *supra*] (indicated in the boxes in Figure 5) included a 27kD amino terminal
- 10 fragment, a 46kD collagen binding fragment, a 75kD cell binding fragment, a 33kD heparin binding fragment associated with a second 66kD fragment (33/66kD fragment), and a 31kD carboxy terminal fragment. Recombinant fragments from the 33kD domain (see Figure 5) were prepared as described elsewhere [Huesbsch *et al.*, *Cir. Res.* **77**:43 (1995); Verfaillie *et al.*, *Blood* **84**:1802 (1994)].

15

Binding assay to immobilized fibronectin, VCAM and fragments. Fibronectin, its recombinant fragments, or rsVCAM were noncovalently immobilized through passive adsorption to 60 well Terasaki trays (wetable polystyrene [plasma treated], Robbins Scientific, Sunnyvale, CA). Briefly, substrates were reconstituted (50

- 20 mg/mg) in phosphate buffered saline (DPBS) and allowed to coat the Teresaki trays for 90 min at 37°C. The polystyrene surface was subsequently blocked with 5% bovine serum albumin (Sigma) for at least 3 hours at 37°C. Prior to use, the plates were washed five times with DPBS and excess liquid was removed from the wells. This procedure was shown to result in the preferred dense, multilayer packing of
- 25 adsorbed fibronectin molecules ($2.7\mu\text{g}/\text{cm}^2$) [van der Flier *et al.*, 1995, *supra*].

Bacteria were labeled with fluorescein isothiocyanate as described [Geelen *et al.*, *Infect. Immun.*, **61**:1538-1543 (1993)] and were resuspended in DPBS supplemented with 0.05% glucose and Ca^{++} and Mg^{++} . Bacterial suspensions were brought to an

- 30 A_{620} of 0.04, previously established to equal 1×10^7 cfu/ml of *S. pneumoniae*. $10\mu\text{l}$

of bacteria were added to each well and incubated statically for 1 hour at 37°C. Unbound bacteria were eliminated by washing five times with DPBS. Bound bacteria were fixed to the surface by incubation with 2.5% glutaraldehyde solution for 3 minutes. Bacteria were counted visually with an inverted microscope (Nikon) equipped for fluorescence with an IF DM 510 filter. Binding was expressed as the number of attached bacteria per 0.25mm² surface area. Values were corrected for non-specific binding by subtracting adherence to uncoated wells and are presented as the mean \pm standard deviation of at least 3 experiments with 3 to 6 wells/plate/experiment. For some experiments, binding was performed in the presence of rabbit polyclonal anti-pneumococcus R6 antibody or one of the following monoclonal antibodies: anti-choline (TEPC-15), anti-PspA Xi126 and XiR278 [McDaniel *et al.*, *Microb. Pathog.*, 13:261-9 (1992)]; anti-pneumococcal surface adhesin A PsaA [Sampson *et al.*, *Infect. Immun.*, 62:319-324 (1994)]; anti-VLA-4 (R & D Systems); and anti-VCAM-1 (R & D Systems). Alternatively, potential inhibition of pneumococcal binding was assessed in the presence of purified recombinant PspA or PsaA.

EXAMPLE 5: THE 112KD CHOLINE BINDING PROTEIN, CbpA MEDIATES PNEUMOCOCCAL ADHESION TO HUMAN CELLS

In this Example, the novel 112 kDa CBP is further characterized. It functions as an adhesin, mediating adherence to cytokine activated human cells and participates in pneumococcal colonization of the nasopharynx, and is a determinant of virulence.

Using choline immobilized on a solid matrix, a family of CBPs was purified from a *pspA*⁻ strain of pneumococcus. Polyclonal antibody to these CBPs passively protected mice infected intraperitoneally with a lethal challenge of pneumococci. The predominant member of this family of proteins, CbpA (or CBP-112), is a 112 kDa surface exposed protein that reacts with human convalescent sera. Sequence analysis of the corresponding gene shows a unique N-terminal sequence and a C-

terminal domain comprised of 10 repeated choline binding domains similar to PspA. A *cbpA* defective mutant shows a greater than 50% reduction in adherence to cytokine activated human cells and failed to bind to immobilized sialic acid or laco-N-neotetraose. This mutant also shows a 100-fold reduction in virulence in an animal model for nasopharyngeal colonization. There is no difference between the parent strain and this mutant to an intraperitoneal model for sepsis. This data for CbpA extends the important functions of the CBP family to bacterial adherence and virulence and represents a new pneumococcal vaccine candidate.

Material and Methods

Strains and media. Serotype 2 *S. pneumoniae* strain D39, R6x (uncapsulated derivative of D39) strain SIII (capsular serotype 3) (each obtained from The Rockefeller University collection). Strain P317 is a 6B capsular serotype. A *lytA* deficient strain has been described [Tomasz, 1988]. LM91 was reported previously (Larry McDaniel, University of Alabama, Birmingham, AL). All strains were plated on tryptic soy agar supplemented with sheep blood (TSAB) to a final concentration of 3% (v/v). Cultures were grown without aeration at 37°C in 5% CO₂ in a liquid semi-synthetic casein hydrolysate medium supplemented with yeast extract (C+Y medium), as described in Example 1. Pneumococci with integrated plasmids were grown in the presence of 1 mg ml⁻¹ erythromycin. Colony morphology was assessed on tryptic soy agar onto which 5000 U of catalase (Worthington Biochemical Co., Freehold, NJ) were added.

Isolation of CBPs. To prepare the immobilized choline affinity matrix, vinylsulfone-activated agarose beads (1 g; Sigma, St. Louis, MO) were washed twice with 10 ml of distilled water and then rotated overnight at room temperature in 10 mM choline in sodium carbonate buffer, pH 11.5. The beads were again washed with an additional 10 ml PBS to remove unbound choline.

- The choline-agarose beads were added to a 400 ml culture of pneumococci (10^8 cfu/ml) and incubated for 30 min. The bacteria-bead complex were pelleted by centrifugation at $8000 \times g$ for 15 min, and resuspended in 50 ml C+Y medium. Bacteria were lysed with Triton X-100 (0.5%; rotation for 20 min) in the presence
- 5 of leupeptin (20 mg ml^{-1}) and phenyl-methyl-sulfonyl fluoride (100 mg ml^{-1}). The lysate was centrifuged at $1000 \times g$ for 5 min to pellet the beads. The beads were then washed with three volumes (30 ml) of phosphate buffered saline (PBS) adjusted to 0.5M NaCl, to remove any non specific bound material. Specific choline bound material was eluted with PBS adjusted to a final concentration of 10% choline (w/v).
- 10 This eluate was dialyzed against PBS and then passed through a Centricon 10 ultrafilter (Amicon Inc. Beverly, MA). The retentate contained the CBPs.

- Analytical and Immunological Methods.* The CBPs were analyzed by 7.5% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and by Western
- 15 blotting after electrophoretic transfer to an Immobilon-P transfer membrane (Millipore Corporation, Bedford, MA). For some experiments, pneumococci were labelled with fluorescein isothiocyanate (FITC) as described in the adherence assay. Western blot analysis was carried out with anti-FITC (Molecular Probes), anti-CBP or covalent antisera each at a dilution of 1:1000. Bound antibody was detected
- 20 with peroxidase conjugated goat anti-rabbit serum detected with the Chemiluminesce Kit from Amersham (Cambridge, MA). The convalescent sera was a mixture of sera pooled from five patients collected one month after recovery from bacteremia and pneumonia. The capsular serotypes of these infecting pneumococci were 3, 7, 14, 22, and 23.

- 25
- For Western analysis of CBPs in phenotypic variants, opaque and transparent variants derived from strain R6 were grown to equal density in C+Y medium (O.D.₆₂₀ of 0.4). The cells were washed in PBS and resuspended in 1/20th the original culture volume in PBS with or without 2% choline chloride. The cells
- 30 were incubated for 15 min. At 4°C and the cell and supernatant (eluate) fractions

separated. Proteins in the eluate were separated on 10% SDS-PAGE gels, transferred to Immobilon-P, and probed with the antiserum to CBPs.

- To obtain primary structural information from individual CBPs, concentrated
- 5 samples from the choline affinity column were applied to SDS polyacrylamide gel and the separated proteins transferred by electrophoresis to an Immobilon-P^{sq} transfer membrane. Proteins were identified following staining with 0.1% amido black : 10 % acetic acid : 40 % methanol. Membranes containing individually stained protein bands were excised and submitted to the Rockefeller University
- 10 Protein Sequencing Center for either N-terminal or internal sequence analysis.

Preparation of rabbit anti-CBP antiserum. Antiserum was generated by HRP Inc (Denver, PA), according to the methods described in Example 1.

- 15 *Cloning and sequencing of cbpA.* DNA techniques including plasmid preparations, restriction endonuclease digests, ligations, transformations into *E. coli* and gel electrophoresis were according to standard procedures known in the art.

- Initial DNA sequence information for cbpA (SEQ ID NO: 24) was obtained from
- 20 DNA fragments generated by anchored PCR. These fragments were generated using degenerate primers derived from the partial primary structure of internal fragments of the purified protein (SEQ ID NO: 25) purified by SDS polyacrylamide gel electrophoresis. Further sequence information was obtained from PCR generated fragments produced with templates to the end regions of the known
- 25 fragments and the anchored universal primer. The polymerase chain reaction (PCR) was performed using the Gene Amp Kit (Perkin Elmer Cetus). DNA sequencing was performed on PCR fragments with an ABI373 DNA sequencer using dye labeled dideoxy terminator chemistry.

A degenerate oligonucleotide corresponding to the amino-terminal amino acid sequence (SEQ ID NO: 1) of CbpA was prepared (5'GCTCTTNCTCGATGTCTCNGTNGCCAT3'; sense) (SEQ ID NO: 22). The antisense oligonucleotide (AGCATAGTCTTCTTCGACTTGTTGATCATC) (SEQ ID NO: 23) was prepared according to the internal amino acid sequence (SEQ ID NO: 10) that was similar to an internal sequence of PspA. LM91 total chromosomal DNA was prepared as described previously and was added (20 ng) to a PCR reaction mixture containing 1 mM each of the sense and antisense primers described above, according to the recommendations provided with AmpliTaq DNA polymerase. Thirty cycles of PCR were performed with oligonucleotide hybridization at 55°C. The DNA band at approximately 600 bp was isolated following agarose gel electrophoresis with an ultrafree-MC filter unit (0.45 mm, Millipore Corporation, Bedford, MA), digested with *Sau3A* and ligated in the *Bam*HI site of the vector pJDC9. This mixture was transformed into *E. coli* DH5 α and a single recombinant clone that contained the vector with the insert was identified.

Further sequence information was obtained from fragments generated by 'anchored' PCR using primers based on the ends of the known fragments and the universal primer anchored to restriction enzyme fragments of chromosomal DNA. Nucleotide sequencing was performed with an ABI 3373 DNA sequencer using dye labeled dideoxy terminator chemistry.

A *cbpA* deficient mutant was created by insertion duplication mutagenesis. First, a 643 base pair DNA fragment corresponding to nucleotide 554 through 1196 of *cbpA* was generated by the PCR. This fragment encompasses the coding region for amino acids 155 through 372. A *Sau*III digest of this fragment produced two fragments of 201 and 195 base pair which were isolated from an agarose gel, ligated into the *Bam*HI site of pJDC9, and transformed into *E. coli* (DH5 α). A single transformant that was transformed into the D39 or 6B strains of

- pneumococcus and two transformants designated SPRU625 (D39) and SPRU632 (6B) were identified that did not express CbpA by Western analysis. Southern analysis confirmed chromosomal integration of the vector disrupting *cbpA* in SPRU625 (Data not shown). Chromosomal DNA digested with either *Hind*III or
- 5 *Eco*RV was separated by electrophoresis and, transferred bi directionally to Hybond-N (Amersham). The membranes were probed with 643 base pair PCR fragment labeled with horseradish peroxidase (Amersham RPN 3000) as recommended by the manufacturer.
 - 10 Southern analysis was carried out on electrophoretically separated chromosomal DNA digested with *Hind*III and *Eco*RV, transferred bidirectionally to Hybond-N (Amersham) and probed with the 600 bp PCR fragment labelled with horseradish peroxidase (Amersham RPN 3000) as recommended by the manufacturer.
 - 15 *Bacterial adhesion assay.* The human Type-II lung cell line A549 (American Type Culture Collection) was cultured in Nutrient mixture F12 Ham medium (Sigma, St. Louis, MO) supplemented with 10 % fetal calf serum (Sigma). Human umbilical vein endothelial cells (HUVEC; Clonetics, San Diego, CA) were cultured in medium M199 (Sigma). At confluence, the cells were prepared for subculture with
 - 20 trypsin-0.05 % EDTA (Sigma). For adherence assays, cells were transferred to Terasaki 60-well culture dishes (Robbins Scientific, Sunnyvale, CA) and cultured for another 24 to 48 hours to form a confluent monolayer. To activate the human cells, some monolayers were incubated with TNF α (10ng/ml for 2 hours; Boehringer-Manheim) or IL-1 β (5 ng/ml for 4 hours; Sigma). Prior to the
 - 25 adherence assay, culture fluid was removed by washing the monolayers twice with tissue culture medium.
- Bacteria at an OD₆₂₀ of 0.4 or 0.6 were washed in 1 ml carbonate buffer (0.05 M sodium carbonate, 0.1 M sodium chloride), resuspended in the same buffer and
- 30 labeled with FITC (Sigma; 1 mg ml⁻¹) for 20 min in the dark at room temperature

(LO). After 3 washes with carbonate buffer, the bacteria were resuspended in medium M199 without antibiotics and 5×10^7 pneumococci were incubated per well for 30 min at 37°C in the presence of 5% CO₂. After the removal of unbound bacteria by washing the monolayers five times with M199, the cells and bacteria were fixed in 2.5% glutaraldehyde for 3 min and washed five times with PBS. Adherent pneumococci were counted visually with an inverted microscope (Diaphot-TDM; Nikon Inc., Melville, NY) equipped for epifluorescence with an IF DM-510 filter and expressed as the number of attached bacteria per 100 lung cells. Values for 6-9 wells were averaged and each experiment was performed 3-6 times.

10

To test the ability of the mixture of CBPs to affect adherence to eucaryotic cells, the assay was modified such that monolayers were plated in 96 well dishes (Falcon) coated with 0.2% gelatin and at confluence were incubated with a range of concentrations of the mixture of CBPs ($1 \mu\text{g}$ to 1 mg ml^{-1}) for 15 min. After washing, the CBP-treated monolayers were challenged with 5×10^6 pneumococci for 30 min, washed and adherence was quantitated as fluorescence intensity measured in a Cytofluor II (Perseptive) with excitation at 485 nm and emission at 530 nm.

15

Adherence to glycoconjugates was assessed by coating Terasaki plates overnight with 100 M of 6'sialyllactose-HSA, lacto-N-neotetraose-HSA, N-acetylglucosamine- β 1,4-glucose-HSA or N-acetylglucosamine- β 1,3-glucose-HSA (Neose Inc., Horsham, PA). Wells were washed and 1×10^7 FITC labelled pneumococci were added for 30 min at 37°C. Unbound cells were washed away three times with PBS and adherence was quantitated visually as described above. Each glycoconjugate was tested in 18 wells during three experiments.

25

Passive protection against systemic challenge. Outbred CF1 mice were housed under specific pathogen free conditions in accordance with institutional and NIH guidelines. Encapsulated pneumococci were grown for 5 hours in C+Y medium

30

and diluted in PBS. Two groups of ten mice received an inoculum of 4.5×10^4 cfu of D39 by injection into the peritoneal cavity. One hour after bacterial challenge, one group of mice received rabbit-anti-CBP serum intraperitoneally (0.5 ml 1:10 in PBS). Control animals received pre-immune serum. A second experiment was

5 performed at a higher challenge dose, with two groups of five mice each receiving 8.4×10^4 cfu of pneumococci. For challenge with SIII, groups of five mice received 200 cfu intraperitoneally of untreated SIII or SIII preincubated for 30 min with anti-CBP antiserum. SIII is a highly virulent strain with an LD_{50} of less than 10 bacteria.

10

Nasopharyngeal challenge. Nasopharyngeal colonization of 1 to 5-day old Sprague-Dawley rats by pneumococci was carried out as described previously using isolates of two capsular serotypes 2 (D39) and 6B (P317). For each experiment, litters were randomized and sorted into groups ten pups per strain of pneumococci. Each

15 pup received equal intranasal inocula of 10^4 cfu in 10 ml of PBS of either the parent strain (D39 or P317) or the isogenic mutant containing a defined mutation in *cbpA*. As an additional control an isogenic strain (P354) which contains a disruption in the surface IgA1 protease gene, *iga*, was used. Colonization was assessed at 24 and 120 hours post-inoculation. To insure accurate evaluation of recovered bacteria, the

20 fluid from the nasal washes were diluted in series, plated and colony counts determined. Results are expressed as the geometric mean of each group \pm the standard deviation ($n=20$).

Results

25 *Characterization of CBPs.* As in Example 1, a mixture of CBPs was prepared by incubating a choline affinity matrix with a *pspA* deficient strain of bacteria, followed by lysis of cells, treating the beads with a high salt solution to remove non-specifically bound material and eluting the CBPs with a 10% choline solution. The CBP, LytA (muramidase), was tracked through the preparative and analytical

30 procedures by western analysis and served as a positive control. Electrophoretic

analysis of the CBP preparation showed at least eight proteins larger than 45 kDa. A protein with an apparent molecular mass of 112 kDa was the most abundant. The muramidase, LytA, is present in the preparation with a molecular mass of 36 kDa, confirming specificity of the technique.

5

To determine whether this mixture of CBPs contained protective antigens, polyclonal antibody was raised to the CBP mixture and tested for ability to protect mice passively from intraperitoneal challenge with bacteria at a concentration 1000-fold greater than the LD₅₀ (Figure 16A). Nine of ten animals receiving pre-immune serum died over 3 days in contrast with animals receiving anti-CBP antiserum. In a second experiment using a 2-fold higher inoculum, 100% of the pre-immune treated animals died over one day while all anti-CBP-treated animals survived six days. The antiserum was then tested for protection against sepsis induced by the heterologous capsule type (Type 3). All control mice receiving intraperitoneal SIII alone or SIII with pre-immune serum were bacteremic at 24 hours with a wide range of bacterial densities from 2×10^4 to 1×10^7 cfu/ml. In contrast, animals receiving SII with anti-CBP serum or anti-Type III serum showed less than 10^4 bacteria/ml of blood. This difference was reflected in survival at 36 and 48 hours. Half of control animals died by 48 hours while none of the anti-CBP or anti-Type

10 serum died over 3 days in contrast with animals receiving anti-CBP antiserum. In a second experiment using a 2-fold higher inoculum, 100% of the pre-immune treated animals died over one day while all anti-CBP-treated animals survived six days.

15 The antiserum was then tested for protection against sepsis induced by the heterologous capsule type (Type 3). All control mice receiving intraperitoneal SIII alone or SIII with pre-immune serum were bacteremic at 24 hours with a wide range of bacterial densities from 2×10^4 to 1×10^7 cfu/ml. In contrast, animals receiving SII with anti-CBP serum or anti-Type III serum showed less than 10^4 bacteria/ml of blood. This difference was reflected in survival at 36 and 48 hours. Half of control animals died by 48 hours while none of the anti-CBP or anti-Type

20 III antibody died during the same time interval. Protection was eventually overcome and all animals died by 72 hours regardless of treatment.

To determine if the CBP mixture might contain adhesins, the CBP mixture was tested for its ability to inhibit the adherence of pneumococcus to endothelial and epithelial cells. Pre-incubation of the eukaryotic cells with the CBPs resulted in a decrease in adherence to epithelial cells by about 50% and to endothelial cells by greater than 80% (Figure 17A). This phenomenon was dose dependent with half maximal activity requiring approximately 60 μ g/ml of CBP (Figure 17B).

25 epithelial cells. Pre-incubation of the eukaryotic cells with the CBPs resulted in a decrease in adherence to epithelial cells by about 50% and to endothelial cells by greater than 80% (Figure 17A). This phenomenon was dose dependent with half maximal activity requiring approximately 60 μ g/ml of CBP (Figure 17B).

Based on the indications that the mixture of CBPs contained potential protective antigens and adhesive ligands, individual CBPs were studied further.

Characteristics were sought which would be important for bioactive CBPs, specifically localization to the pneumococcal surface and reactivity with human

5 convalescent antisera and the protective anti-CBP serum. Intact pneumococci were labelled with FITC and then subjected to CBP fractionation. SDS-PAGE and western blot analysis using anti-FITC antibody showed four proteins labelled with FITC in both whole cell preparations and CBP preparations. This is consistent with access of this group of CBPs to the pneumococcal extracellular milieu in the native
10 bacteria. Five CBPs reacted with human convalescent antisera and the 112 kDa band was prominently labelled in preparations from whole pneumococci. This same pattern occurred using the anti-CBP antiserum, which prominently labelled the 112 kDa band in intact pneumococci and CBP preparations.

15 As a result of localization of the 112 kDa band to the pneumococcal surface and its strong reactivity with convalescent antisera and protective rabbit anti-CBP-sera prepared from whole bacteria or the described CBP fractionation procedure, it was determined that the 112 kDa protein is major protein component of the CBP mixture and thus designated CbpA.

20

Cloning and sequence analysis of cbpA. Direct N-terminal amino acid analysis of CbpA revealed a unique sequence (SEQ ID NO:1) with no similarity to proteins in any published data base. In contrast, an internal sequence (SEQ ID NO:10) was identical to an internal sequence of the CBP, PspA. Since a *pspA* deficient strain
25 was used to identify *cbpA*, this result was not an artifact due to a genetic rearrangement of *pspA* but probably represented a gene composed of a unique 5' coding region and a 3' coding region similar to *pspA*. Using a PCR based strategy, *cbpA*, was cloned and sequenced. Southern analysis using a probe unique to the 5' region (see Experimental procedure section) indicated only one copy of *cbpA* in the
30 LM91 chromosome (data not shown).

Analysis of the derived protein sequence of *cbpA* brought to light several unique properties. CpbA is a protein of 630 aa with an apparent molecular mass of 71 kDa. This is different from the calculated molecular mass (112 kDa) based on its migration properties by SDS-PAGE. The presence of a standard N-terminal signal sequence implies SecA dependant export. The protein has two distinct domains. The N-terminal region (1-340) has no obvious similarities to proteins entered into published databases. Ganier-Robson analysis predicted six alpha-helical regions from residues 10-350. Five coiled coil structures greater than 50 aa in length in this same region were predicted with the paircoil algorithm. In contrast, the C-terminal region (341-360) is nearly identical (> 95%) to PspA. This similarity extends to the corresponding nucleotide sequence. Common features are a proline rich region (340-370) and ten tandem, direct repetitive sequences of 20 aa (381-584) that represent the choline binding motif that is the signature of the choline binding proteins.

Phenotypic variation and expression of the choline binding proteins. Since opaque and transparent variants of pneumococci were shown to have differences in the expression of the CBP LytA, the expression of the CBPs were compared between opaque and transparent variants of a noncapsulated strain (R6x) with anti-CBP serum. All though all bands were present in both phenotypes, several bands differed between the opaque and transparent cell types. Transparent organisms had significantly higher amounts of LytA, as previously documented. In addition, the transparent variant expressed increased amounts of CbpA and PspA. In contrast, the opaque pneumococci had more PspA. In these data minor bands between 70 and 90 kD appeared to be degradation products of PspA and CbpA. There was a protein of approximately 50 kDa present in the transparent organisms regardless of the presence of choline suggesting that this was not a CBP. As an added control, the identity of these CBPs, LytA and PspA, was confirmed by the absence of these bands in material obtained from mutants with defects in the corresponding genes (data not shown).

Functional analysis of CbpA- mutant. To assess bio activities that might be affected by CbpA, genetic mutants were constructed in a variety of assays.

- 5 The growth rates of two CbpA⁻ mutants in semisynthetic medium was equivalent to wild type and the rate of phenotypic variation from transparent to opaque colonial variants was not altered from the parental strain (data not shown). Other physiological properties of pneumococci were also unchanged, including efficiency of DNA transformation of a streptomycin resistance marker, lysis in stationary phase, and penicillin-induced lysis (data not shown).
- 10 The capsular serotype 2 *cbpA* deficient mutant was tested for the ability to adhere to human lung epithelial cells and endothelial cells and glycoconjugates. The mutant adhered to about 80% of parental strain levels (Figure 18A, B).
- 15 This was consistent with the equivalent ability of the two strains to adhere to immobilized GalNAc- β 1,4-Gal and GalNAc- β 1,3-Gal, sugars known to be receptors on these eucaryotic cells (Cundell) (Figure 18C). A difference in adherence between the parent and mutant was detected, however, with cytokine activated cells. Adherence of the parental strain to cytokine activated cells increased to ~135% of
- 20 resting cell values. In contrast, adherence of the *cbpA* deficient mutant failed to increase for either lung or epithelial cells (Figure 18A). This defect in adherence to cytokine activated cells was consistent with the absence of adherence of the mutant to purified glycoconjugates (sialic acid and lacto-N-neotetraose, LnNT) known to be receptors for pneumococci on cytokine activated cells (Figure 18C).
- 25 An infant rat model of pneumococcal carriage was used to determine if the ability of CbpA to mediate adherence to human epithelial cells correlated with the ability of the pneumococcus to colonize the nasopharynx. In this animal model, the pneumococcus persists on the mucosal surface at high levels for many days
- 30 following a small intranasal inoculation. In contrast to the controls which were a

10

15

20

25

30

and vascular endothelial cells. These cells are *in vitro* models for pneumococcal adherence at sites of pneumococcal infection (NEJM). This suggested that the CBPs bound to receptors on the human cells and thereby prevented CBP mediated bacterial attachment. The demonstration of a direct CBP human cell interaction that leads to a loss of adherence distinguishes the CBPs from all other pneumococcal proteins that have been identified in genetic studies as affecting bacterial adherence. While loss of these other elements decreases the capacity of the bacteria to adhere, there is no indication that they exhibit competitive inhibition of adherence, a key characteristic of a structural adhesin. These adherence and virulence properties of the CBP mixture encouraged further analysis of individual CBPs.

CbpA was chosen for further studies because it was the most abundant CBP in the mixture, it was surface exposed and reacted strongly with both human convalescent antibody and the mouse protective anti-CBP serum, both on whole bacteria and after purification in a CBP preparation. Although the sequence predicts a molecular weight of 71 kDa, the protein migrates with an apparent mass of 112 kDa on SDS-PAGE, a discrepancy which is also characteristic of another CBP, PspA. The choline binding domains of CbpA appears to be identical to those of PspA both at the amino acid and nucleotide level and thus chimeric gene structures for these elements. The N-terminal domains of the two CBPs are very different in primary sequence, yet they share common structural features such as several large alpha-helical regions and five coiled coil regions. This would suggest that CbpA may be a fibrous protein analogous of PspA, tropomyosin, troponin, and the M protein family of streptococcal proteins. It seems reasonable that the N-terminal domain of CbpA contains one or two lectin domains capable of binding LnNT and sialic acid. Such lectin domains would be expected to remain conserved between strains and serotypes since they must retain the ability to bind the target humans glycoconjugates receptors in order to retain virulence. This conservation may contribute to cross protection between various serotypes such as was observed for the anti-CBP antiserum.

CbpA appears to vary in expression coincident with the phase transition between the opaque and transparent variants. Some CBPs, such as PspA, are expressed in higher amounts in opaque variants. CbpA varies in the same manner as LytA such that transparent variants express increased amount of this protein. The significance of increased PspA in opaque organisms and the function of this CBP is unknown. In contrast the increased expression of LytA and CbpA by transparent forms correlates with increased adherence and colonization. A role for LytA in colonization could not be demonstrated. In contrast, the role of CbpA in adherence and colonization was confirmed directly by analysis of the *cbpA* deficient mutant.

10

A two step model for pneumococcal adherence has been proposed. Pneumococci initially target a niche in the host by binding surface glycoconjugates, such as GalNAc- β 1,4-Gal and GalNAc- β 1,3-Gal, on the surfaces of eucaryotic cells. A *cbpA* deficient mutant was unaffected in its adherence properties to these cells or the corresponding glycoconjugates. A second adhesive step is believed to lead to invasion and is observed upon cytokine activation of the human cell. New sugar specificities appear on the activated cells resulting in increased pneumococcal adherence. This second level of adherence is distinguished by the ability of sialylated glycoconjugates (6'SL) and LnNT to inhibit bacterial attachment. These sugars are not receptor analogs for resting cells. A *cbpA* deficient mutant lost the ability to bind to cytokine-activated cells or to immobilized 6'SL and LnNT.

20

Together with the ability of purified CBPs to inhibit pneumococcal attachment, these results suggest that CbpA may be a structural adhesin with potential lectin activity. Cytokine activated cells are suggested to express the platelet activating factor (PAF) receptor which can bind the phosphorylcholine of the pneumococcal cell wall. The presence of CBPs attached to the phosphorylcholine would presumably mask phosphorylcholine binding to the PAF receptor. The relative number of free versus CBP-bound phosphorylcholine determinants on the pneumococcal surface is unknown. It appears that both CbpA and phosphorylcholine participate in pneumococcal binding to activated human cells.

30

- CbpA appears to represent the first example of a protein adhesin on the pneumococcal surface. It may act as a bridge between cells wall phosphoorylcholine, bound by the C-terminal choline binding repeats, and human
- 5 cell glycoconjugates, presumably via the N-terminal domain. The binding capacity is restricted to activated human cells and may, therefore, be important in advancing the course of disease from benign colonization to invasion. This is also consistent with results suggesting CbpA is subject to phase variation and that it is a marker for the transparent, colonization proficient, phase variants of pneumococci.
- 10 Accordingly, CBPs, and CbpA (CBP-112) in particular, are attractive targets for vaccine development and passive immunotherapy.
- 15 While the invention has been described and illustrated herein by references to various specific material, procedures and examples, it is understood that the invention is not restricted to the particular material combinations of material, and procedures selected for that purpose. Numerous variations of such details can be implied as will be appreciated by those skilled in the art.

Various references are cited throughout this specification, each of which is incorporated herein by reference in its entirety.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Masure, H. Robert
Rosenow, Carsten I.
Tuomanen, Elaine
Wizemann, Theresa M.

(ii) TITLE OF INVENTION: CHOLINE BINDING PROTEINS FOR
ANTI-PNEUMOCOCCAL VACCINES

(iii) NUMBER OF SEQUENCES: 25

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: David A. Jackson, Esq.
(B) STREET: 411 Hackensack Ave, Continental Plaza, 4th
Floor
(C) CITY: Hackensack
(D) STATE: New Jersey
(E) COUNTRY: USA
(F) ZIP: 07601

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Jackson Esq., David A.
(B) REGISTRATION NUMBER: 26,742
(C) REFERENCE/DOCKET NUMBER: 600-1-158 ..

(ix) TELECOMMUNICATION INFORMATION:

(B) TELEFAX: 201-343-1684

[illegible]

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ala	Arg	Glu	Phe	Ser	Leu	Glu	Lys	Thr	Arg
1				5					10

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP84

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Xaa	Arg	Glu	Phe	Ser	Leu	Glu	Lys	Thr	Arg	Asn	Ile	Gly	Ile	Met	Ala
1				5				10					15		

His	Val	Asp	Ala	Gly	Lys	Thr
						20

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (A) DESCRIPTION: cBP80

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Xaa	Lys	Xaa	Xaa	Trp	Gln	Xaa	Lys	Gln	Tyr	Leu	Lys	Glu	Asp	Gly	Ser
1				5					10					15	
Gln	Ala	Ala	Asn	Glu	Xaa	Val	Phe	Asp	Thr	Ala					
			20					25							

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (A) DESCRIPTION: cBP78

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Gln	Lys	Ile	Ile	Gly	Ile	Asp	Leu	Gly	Thr	Thr	Asn	Ser	Ala	Val	Ala
1				5					10					15	
Val	Leu	Glu	Gly	Thr	Glu	Ser	Lys	Ile	Ile	Ala	Asn	Pro	Glu		
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP70

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Xaa	Xaa	Xaa	Glu	Val	Ala	Lys	Xaa	Ser	Gln	Asp	Thr	Thr	Thr	Ala	Ser
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP60

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Xaa	Asn	Glu	Arg	Val	Lys	Ile	Val	Ala	Thr	Leu	Gly	Pro	Ala	Val	Glu
1					5					10				15	

Gly Arg Gly

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: cBP50 pep

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Xaa	Ile	Ile	Xaa	Xaa	Val	Tyr	Ala	Arg	Glu	Val	Leu	Asp	Ser	Arg	Gly
1				5					10					15	

Asn Pro

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: cBP112-Int1

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Glu	Asp	Arg	Arg	Asn	Tyr	Pro	Thr	Asn	Thr	Tyr	Lys
1				5						10	

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP112-Int2

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Xaa	Asp	Asp	Gln	Gln	Ala	Glu	Glu	Asp	Tyr	Ala
1				5					10	

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Trp Gln Pro Pro Arg Ala Arg Ile
1 5

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: 50 KDa protein

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *S. pneumoniae*

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ile Thr Asp Val Tyr Ala Arg Glu Val Leu Asp Ser Arg Gly Asn Pro
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (A) DESCRIPTION: 50 KDa protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: B. subtilis

(xi) SEQUENCE DESCRIPTION: SEQ ID,NO:13:

Ile	Val	Asp	Val	Arg	Ala	Arg	Glu	Val	Leu	Asp	Ser	Arg	Gly	Asn	Pro
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 210 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: S. pneumoniae

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TCCCGCGGCA ACCCAACACT TGAAGTAGAA GTTTACACTG AATCAGGTGC TTTCGGACGT

GGTATGGTTC CATCAGGAGC TTCTACTGGT GAACACGAAG CAGTTGAACT TCGCGACGGT 120
 GACAAATCTC GTTACGGTGG TCTTGGTACA CAAAAGCTG TTGACAACGT AAACAACATC 180
 ATTGCTGAGG CCATCATTGG CTACGATGTA 210

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: B. subtilis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TCCCGCGGCA ACCCAACAGT TGAAGTTGAA GTATATACAG AAACAGGAGC TTTCGGCCGC 60
 GCATTAGTGC CAAGCGGAGC TTCTACAGGT CAATACGAAG CGGTTGAGCT TCGTGACGGC 120
 GACAAAGACC GTTACCTTGG AAAAGGCGTG TTAACAGCTG TTAACAACGT AAACGAAATC 180
 ATTGCTCCAG AGCTTCTTGG CTTTGATGTA 210

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 amino acids
- (B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID.NO:16:

Ser	Arg	Gly	Asn	Pro	Thr	Leu	Glu	Val	Glu	Val	Tyr	Thr	Glu	Ser	Gly
1				5					10					15	

Ala	Phe	Gly	Arg	Gly	Met	Val	Pro	Ser	Gly	Ala	Ser	Thr	Gly	Glu	His
			20					25					30		

Glu	Ala	Val	Glu	Leu	Arg	Asp	Gly	Asp	Lys	Ser	Arg	Tyr	Gly	Gly	Leu
		35					40						45		

Gly	Thr	Gln	Lys	Ala	Val	Asp	Asn	Val	Asn	Asn	Ile	Ile	Ala	Glu	Ala
	50						55				60				

Ile	Ile	Gly	Tyr	Asp	Val
65				70	

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *B. subtilis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Ser Arg Gly Asn Pro Thr Val Glu Val Glu Val Tyr Thr Glu Thr Gly
1 5 10 15

Ala Phe Gly Arg Ala Leu Val Pro Ser Gly Ala Ser Thr Gly Gln Tyr
 20 25 30

Glu Ala Val Glu Leu Arg Asp Gly Asp Lys Asp Arg Tyr Leu Gly Lys
 35 40 45

Gly Val Leu Thr Ala Val Asn Asn Val Asn Glu Ile Ile Ala Pro Glu
 50 55 60

Leu Leu Gly Phe Asp Val
65 70

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1511 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TCCCGCGGCA ACCCAACACT TGAAGTAGAA GTTTACACTG AATCAGGTGC TTTCGGACGT	60
GGTATGGTTC CATCAGGAGC TTCTACTGGT GAACACGAAG CAGTTGAACT TCGCGACGGT	120
GACAAATCTC GTTACGGTGG TCTTGGTACA CAAAAAGCTG TTGACAACGT AAACAACATC	180
ATTGCTGATT CTATCATTGG CTACGATGTA CGTGATCAAC AAGCTATTGA CCGTGCTATG	240
ATCGCACTTG ACGGTACTCC TAACAAAGGT AAATTGGGTG CGAATGCAAT CCTCGGTGTG	300
TCTATCGCTG TAGCTCGTGC TGCTGCTGAC TACCTTGAAA TCCCACTTTA 'CAGCTACCTT	360
GGTGGATTCA AACTAAAGT TCTTCCAACCT CCAATGATGA ACATCATCAA CGGTGGTTCT	420
CACTCTGACG CTCCAATCGC TTTCCAAGAG TTCATGATCT TGCCAGTTGG TGCGCCAACA	480
TTTAAAGAAA CCCTTCGTTA CGGTGCTGAA ATCTTCCACG CTCTTAAGAA AATCCTTAAA	540
TCACGTGGTT TGGAAGTGC CGTAGGTGAC GAAGGTGGAT TCGCTCCTCG TTTCGAAGGA	600
ACTGAAGATG GTGTTGAAAC TATCCTTGCT GCGATTGAAG CTGCTGGATA TGTACCAGGT	660
AAAGACGTAT TTCTCGGATT TGACTGTGCT TCATCAGAAT TCTACGATAA AGAACGTAAA	720
GTTTACGACT AACTAAATT TGAAGGCGAA GGTGCTGCTG TTCGTACATC TGCAGAACAA	780
ATCGACTACC TTGAAGAATT GGTTAACAAA TACCCAATCA TCACTATTGA AGATGGTATG	840
GATGAAAACG ACTGGGATGG TTGGAAAGCT CTTACTGAAC GTCTTGGTAA GAAAGTACAA	900
CTTGTTGGTG ACGACTTCTT CGTAACAAAC ACTGACTACC TTGCACGTGG TATCCAAAAA	960
GGTGCTGCTA ACTCAATCCT TATCAAAGTT AACCAAATCG GTACTCTTAC TGAAACTTTT	1020
GAAGCTATCG AAATGGCTAA AGAAGCTGGT TACTGCTG TTGTATCACA CCGTTCAGGT	1080

GAAACTGAAG ATTCAACAAT CGCTGATATC GCGATTGCAA CTAACGCAGG ACAAATCAAG 1140
 ACTGGTTCAC TTTCACGTAC AGACCGTATC GCTAAATACA ACCAATTGCT TCGTATCGAA 1200
 GACCAACTTG GTGAAGTAGC AGAATATCGT GGATTGAAAT CATTCTACGA CCTTAAAAAA 1260
 TAAAATAGTA CAGTGAAC TA TTTTATCCCC GAACCATGAA ATTCAAATTC CGGCCCTTGA 1320
 TTTAACAAC TTCTAGCCCC CCGATTTATC CGAGGGGATA TTTTGT TTTT AAAGGGCAAA 1380
 AAGGACAAAC TTTTAAAAT CTTTTCATAC ATTACCATAC TTGATGCTCC CCTCCTTCTT 1440
 GTACTGGTTT AAGAATATAC TCCATCGATG ACACCGTCGG ATGTTTAGGA TTATTCGACA 1500
 ACAAGTTGAA T 1511

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 420 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ser Arg Gly Asn Pro Thr Leu Glu Val Glu Val Tyr Thr Glu Ser Gly
 1 5 10 15

Ala Phe Gly Arg Gly Met Val Pro Ser Gly Ala Ser Thr Gly Glu His

20

25

30

Glu Ala Val Glu Leu Arg Asp Gly Asp Lys Ser Arg Tyr Gly Gly Leu
 35 40 45

Gly Thr Gln Lys Ala Val Asp Asn Val Asn Asn Ile Ile Ala Asp Ser
 50 55 60

Ile Ile Gly Tyr Asp Val Arg Asp Gln Gln Ala Ile Asp Arg Ala Met
 65 70 75 80

Ile Ala Leu Asp Gly Thr Pro Asn Lys Gly Lys Leu Gly Ala Asn Ala
 85 90 95

Ile Leu Gly Val Ser Ile Ala Val Ala Arg Ala Ala Ala Asp Tyr Leu
 100 105 110

Glu Ile Pro Leu Tyr Ser Tyr Leu Gly Gly Phe Asn Thr Lys Val Leu
 115 120 125

Pro Thr Pro Met Met Asn Ile Ile Asn Gly Gly Ser His Ser Asp Ala
 130 135 140

Pro Ile Ala Phe Gln Glu Phe Met Ile Leu Pro Val Gly Ala Pro Thr
 145 150 155 160

Phe Lys Glu Thr Leu Arg Tyr Gly Ala Glu Ile Phe His Ala Leu Lys
 165 170 175

Lys Ile Leu Lys Ser Arg Gly Leu Glu Thr Ala Val Gly Asp Glu Gly
 180 185 190

Gly Phe Ala Pro Arg Phe Glu Gly Thr Glu Asp Gly Val Glu Thr Ile
 195 200 205

Leu Ala Ala Ile Glu Ala Ala Gly Tyr Val Pro Gly Lys Asp Val Phe
 210 215 220

Leu Gly Phe Asp Cys Ala Ser Ser Glu Phe Tyr Asp Lys Glu Arg Lys
 225 230 235 240

Val Tyr Asp Tyr Thr Lys Phe Glu Gly Glu Gly Ala Ala Val Arg Thr
 245 250 255

Ser Ala Glu Gln Ile Asp Tyr Leu Glu Glu Leu Val Asn Lys Tyr Pro
 260 265 270

Ile Ile Thr Ile Glu Asp Gly Met Asp Glu Asn Asp Trp Asp Gly Trp
 275 280 285

Lys Ala Leu Thr Glu Arg Leu Gly Lys Lys Val Gln Leu Val Gly Asp
 290 295 300

Asp Phe Phe Val Thr Asn Thr Asp Tyr Leu Ala Arg Gly Ile Gln Lys
 305 310 315 320

Gly Ala Ala Asn Ser Ile Leu Ile Lys Val Asn Gln Ile Gly Thr Leu
 325 330 335

Thr Glu Thr Phe Glu Ala Ile Glu Met Ala Lys Glu Ala Gly Tyr Thr
 340 345 350

Ala Val Val Ser His Arg Ser Gly Glu Thr Glu Asp Ser Thr Ile Ala
 355 360 365

Asp Ile Ala Ile Ala Thr Asn Ala Gly Gln Ile Lys Thr Gly Ser Leu
 370 375 380

Ser Arg Thr Asp Arg Ile Ala Lys Tyr Asn Gln Leu Leu Arg Ile Glu
 385 390 395 400

Asp Gln Leu Gly Glu Val Ala Glu Tyr Arg Gly Leu Lys Ser Phe Tyr
 405 410 415

Asp Leu Lys Lys
 420

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 429 base pairs

(B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

AAAAAAGGTA GCAGAAGCTG AGAAGAAGGT TGAAGAAGCT GAGAAAAAAG CCAAGGATCA	60
AAAAGAAGAA GATCGCCGTA ACTACCCAAC CAATACTTAC AAAACGCTTG ACCTTGAAAT	120
TGCTGAGTCC GATGTGAAAG TTAAAGAAGC GGAGCTTGAA CTAGTAAAAG AGGAAGCTAA	180
GGAACCTCGA GACGAGGAAA AAATTAAGCA AGCAAAAGCG AAAGTTGAGA GTAAAAAAGC	240
TGAGGCTACA AGGTTAGAAA ACATCAAGAC AGATCGTAAA AAAGCAGAAG AAGAAGCTAA	300
ACGAAAAGCA GCAGAAGAAG ATAAAGTTAA AGAAAAACCA GCTGAACAAC CACAACCAGC	360
GCCGGTTACT CAACCAGAAA AACCAGCTCC AAAACCAGAG AAGCCAGCTG AACCAACCAA	420
AGCAGAAAA	429

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 142 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Lys	Lys	Val	Ala	Glu	Ala	Glu	Lys	Lys	Val	Glu	Glu	Ala	Glu	Lys	Lys	1	5	10	15
Ala	Lys	Asp	Gln	Lys	Glu	Glu	Asp	Arg	Arg	Asn	Tyr	Pro	Thr	Asn	Thr	20	25	30	
Tyr	Lys	Thr	Leu	Asp	Leu	Glu	Ile	Ala	Glu	Ser	Asp	Val	Lys	Val	Lys	35	40	45	
Glu	Ala	Glu	Leu	Glu	Leu	Val	Lys	Glu	Glu	Ala	Lys	Glu	Pro	Arg	Asp	50	55	60	
Glu	Glu	Lys	Ile	Lys	Gln	Ala	Lys	Ala	Lys	Val	Glu	Ser	Lys	Lys	Ala	65	70	75	80
Glu	Ala	Thr	Arg	Leu	Glu	Asn	Ile	Lys	Thr	Asp	Arg	Lys	Lys	Ala	Glu	85	90	95	
Glu	Glu	Ala	Lys	Arg	Lys	Ala	Ala	Glu	Glu	Asp	Lys	Val	Lys	Glu	Lys	100	105	110	
Pro	Ala	Glu	Gln	Pro	Gln	Pro	Ala	Pro	Val	Thr	Gln	Pro	Glu	Lys	Pro	115	120	125	
Ala	Pro	Lys	Pro	Glu	Lys	Pro	Ala	Glu	Gln	Pro	Lys	Ala	Glu			130	135	140	

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide primer"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GCTCTTNCTC GATGTCTCNG TNGCCAT

27

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide primer"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AGCATAGTCT TCTTCGACTT GTTGATCATC

30

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2462 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCAAGCAGCC ACTTCTTCTA ATATGGCAAA GACAGAACAT AGGAAAGCTG CTAAACAAGC	60
GTCGATGAAT ATATAGAAAA AATGATGAGG GAGATTCAAC TAGATAGAAG AAAACATACC	120
CAAAATGTCG CCTTAAACAT AAAGTTGAGC GCAATTAAAA CGAAGTATTT GCGTGAATTA	180
AATGTTTTAG AAGAGAAGTC GAAAGATGAG TTGCCGTCAG AAATAAAAGC AAAGTTAGAC	240
GCAGCTTTTG AGAAGTTTAA AAAAGATACA TTGAAACCAG GAGAAAAGGT AGCAGAAGCT	300
AAGAAGAAGG TTGAAGAAGC TAAGAAAAAA GCCGAGGATC AAAAAGAAGA AGATCGTCGT	360
AACTACCCAA CCAATACTTA CAAAACGCTT GAACTTGAAA TTGCTGAGTT CGATGTGAAA	420
GTTAAAGAAG CGGAGCTTGA ACTAGTAAAA GAGGAAGCTA AAGAATCTCG AAACGAGGGC	480
ACAATTAAGC AAGCAAAAGA GAAAGTTGAG AGTAAAAAAG CTGAGGCTAC AAGGTTAGAA	540
AACATCAAGA CAGATCGTAA AAAAGCAGAA GAAGAAGCTA AACGAAAAGC AGATGCTAAG	600
TTGAAGGAAG CTAATGTAGC GACTTCAGAT CAAGGTAAAC CAAAGGGGCG GGCAAAACGA	660
GGAGTTCCTG GAGAGCTAGC AACACCTGAT AAAAAAGAAA ATGATGCGAA GTCTTCAGAT	720

TCTAGCGTAG GTGAAGAAAC TCTTCCAAGC TCATCCCTGA AATCAGGAAA AAAGGTAGCA	780
GAAGCTGAGA AGAAGGTTGA AGAAGCTGAG AAAAAAGCCA AGGATCAAAA AGAAGAAGAT	840
CGCCGTA ACT ACCCAACCAA TACTTACAAA ACGCTTGACC TTGAAATTGC TGAGTCCGAT	900
GTGAAAGTTA AAGAAGCGGA GCTTGAACTA GTAAAAGAGG AAGCTAAGGA ACCTCGAGAC	960
GAGGAAAAAA TTAAGCAAGC AAAAGCGAAA GTTGAGAGTA AAAAAGCTGA GGCTACAAGG	1020
TTAGAAAACA TCAAGACAGA TCGTAAAAAA GCAGAAGAAG AAGCTAAACG AAAAGCAGCA	1080
GAAGAAGATA AAGTTAAAGA AAAACCAGCT GAACAACCAC AACCAGCGCC GGCTACTCAA	1140
CCAGAAAAAC CAGCTCCAAA ACCAGAGAAG CCAGCTGAAC AACCAAAAGC AGAAAAACA	1200
GATGATCAAC AAGCTGAAGA AGACTATGCT CGTAGATCAG AAGAAGAATA TAATCGCTTG	1260
ACTCAACAGC AACCGCCAAA AACTGAAAAA CCAGCACAAC CATCTACTCC AAAACAGGC	1320
TGGAAACAAG AAAACGGTAT GTGGTACTTC TACAATACTG ATGGTTCAAT GGCAACAGGA	1380
TGGCTCCAAA ACAACGGTTC ATGGTACTAT CTAAACGCTA ATGGTGCTAT GGCGACAGGA	1440
TGGCTCCAAA ACAATGGTTC ATGGTACTAT CTAAACGCTA ATGGTTCAAT GGCAACAGGA	1500
TGGCTCCAAA ACAATGGTTC ATGGTACTAC CTAAACGCTA ATGGTGCTAT GGCGACAGGA	1560
TGGCTCCAAT ACAATGGTTC ATGGTACTAC CTAAACAGCA ATGGCGCTAT GGCGACAGGA	1620
TGGCTCCAAT ACAATGGCTC ATGGTACTAC CTCAACGCTA ATGGTGATAT GGCGACAGGA	1680
TGGCTCCAAA ACAACGGTTC ATGGTACTAC CTCAACGCTA ATGGTGATAT GGCGACAGGA	1740
TGGCTCCAAT ACAACGGTTC ATGGTATTAC CTCAACGCTA ATGGTGATAT GGCGACAGGT	1800
TGGGTGAAAG ATGGAGATAC CTGGTACTAT CTTGAAGCAT CAGGTGCTAT GAAAGCAAGC	1860
CAATGGTTCA AAGTATCAGA TAAATGGTAC TATGTCAATG GCTCAGGTGC CCTTGCAGTC	1920
AACACA ACTG TAGATGGCTA TGGAGTCAAT GCCAATGGTG AATGGGTAAA CTAAACCTAA	1980

TATAACTAGT TAATACTGAC TTCCTGTAAG AACTTTTTTAA AGTATTCCCT ACAAATACCA 2040
 TATCCTTTCA GTAGATAATA TACCCTTGTA GGAAGTTTAG ATTAAAAAAT AACTCTGTAA 2100
 TCTCTAGCCG GATTTATAGC GCTAGAGACT ACGGAGTTTT TTTGATGAGG AAAGAATGGC 2160
 GGCATTCAAG AGACTCTTTA AGAGAGTTAC GGGTTTTTAA CTATTAAGCC TTCTCCAATT 2220
 GCAAGAGGCT TCAATCTCTG CTAGGGTGCT AGCTTGCGAA ATGGCTCCAC GGAGTTTGGC 2280
 AGCGCCAGAT GTTCCACGGA GATAGTGAGG AGCGAGGCCG CGGAATTCAC GAACTGCGAC 2340
 GTTTTCTCCT TTGAGGTAA TCAATCGTTT CAAGTGTTTCG TAGGCGATCT TCATCTTGTC 2400
 TTCAAAGGTC AAATCAGGTA GGATTTCTCC TGTTTCAAAG TTTATGGTGG CCCTGGTTGA 2460
 AG 2462

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 631 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met Met Arg Glu Ile Gln Leu Asp Arg Arg Lys His Thr Gln Asn Val
 1 5 10 15

Ala Leu Asn Ile Lys Leu Ser Ala Ile Lys Thr Lys Tyr Leu Arg Glu

20					25					30						
Leu	Asn	Val	Leu	Glu	Glu	Lys	Ser	Lys	Asp	Glu	Leu	Pro	Ser	Glu	Ile	
35						40						45				
Lys	Ala	Lys	Leu	Asp	Ala	Ala	Phe	Glu	Lys	Phe	Lys	Lys	Asp	Thr	Leu	
50						55						60				
Lys	Pro	Gly	Glu	Lys	Val	Ala	Glu	Ala	Lys	Lys	Lys	Val	Glu	Glu	Ala	
65						70						75			80	
Lys	Lys	Lys	Ala	Glu	Asp	Gln	Lys	Glu	Glu	Asp	Arg	Arg	Asn	Tyr	Pro	
			85						90						95	
Thr	Asn	Thr	Tyr	Lys	Thr	Leu	Glu	Leu	Glu	Ile	Ala	Glu	Phe	Asp	Val	
			100						105						110	
Lys	Val	Lys	Glu	Ala	Glu	Leu	Glu	Leu	Val	Lys	Glu	Glu	Ala	Lys	Glu	
115						120						125				
Ser	Arg	Asn	Glu	Gly	Thr	Ile	Lys	Gln	Ala	Lys	Glu	Lys	Val	Glu	Ser	
130						135						140				
Lys	Lys	Ala	Glu	Ala	Thr	Arg	Leu	Glu	Asn	Ile	Lys	Thr	Asp	Arg	Lys	
145						150						155			160	
Lys	Ala	Glu	Glu	Glu	Ala	Lys	Arg	Lys	Ala	Asp	Ala	Lys	Leu	Lys	Glu	
			165						170						175	
Ala	Asn	Val	Ala	Thr	Ser	Asp	Gln	Gly	Lys	Pro	Lys	Gly	Arg	Ala	Lys	
			180						185						190	
Arg	Gly	Val	Pro	Gly	Glu	Leu	Ala	Thr	Pro	Asp	Lys	Lys	Glu	Asn	Asp	
195						200						205				
Ala	Lys	Ser	Ser	Asp	Ser	Ser	Val	Gly	Glu	Glu	Thr	Leu	Pro	Ser	Ser	
210						215						220				
Ser	Leu	Lys	Ser	Gly	Lys	Lys	Val	Ala	Glu	Ala	Glu	Lys	Lys	Val	Glu	
225						230						235			240	

Ala Met Ala Thr Gly Trp Leu Gln Asn Asn Gly Ser Trp Tyr Tyr Leu
450 455 460

Asn Gly Glu Trp Val Asn Glx
625 630

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Masure, H. Robert
Rosenow, Carsten I.
Tuomanen, Elaine
Wizemann, Theresa M.

- (ii) TITLE OF INVENTION: CHOLINE BINDING PROTEINS FOR
ANTI-PNEUMOCOCCAL VACCINES

- (iii) NUMBER OF SEQUENCES: 25

- (iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: David A. Jackson, Esq.
 - (B) STREET: 411 Hackensack Ave, Continental Plaza, 4th
Floor
 - (C) CITY: Hackensack
 - (D) STATE: New Jersey
 - (E) COUNTRY: USA
 - (F) ZIP: 07601

- (v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30

- (vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:

- (viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Jackson Esq., David A.

(B) REGISTRATION NUMBER: 26,742

(C) REFERENCE/DOCKET NUMBER: 600-1-158 ..

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 201-487-5800

(B) TELEFAX: 201-343-1684

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP112

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Xaa	Glu	Asn	Glu	Gly	Ser	Thr	Gln	Ala	Ala	Thr	Ser	Ser	Asn	Met	Ala
1				5					10					15	

Lys	Thr	Glu	His	Arg	Lys	Ala	Ala	Lys	Gln	Val	Val	Asp	Glu
				20				25				30	

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP90

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ala	Arg	Glu	Phe	Ser	Leu	Glu	Lys	Thr	Arg
1				5					10

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP84

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Xaa	Arg	Glu	Phe	Ser	Leu	Glu	Lys	Thr	Arg	Asn	Ile	Gly	Ile	Met	Ala
1					5					10					15

His Val Asp Ala Gly Lys Thr

20

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: cBP80

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Xaa	Lys	Xaa	Xaa	Trp	Gln	Xaa	Lys	Gln	Tyr	Leu	Lys	Glu	Asp	Gly	Ser
1				5				10						15	

Gln	Ala	Ala	Asn	Glu	Xaa	Val	Phe	Asp	Thr	Ala
			20					25		

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (A) DESCRIPTION: cBP78

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Gln	Lys	Ile	Ile	Gly	Ile	Asp	Leu	Gly	Thr	Thr	Asn	Ser	Ala	Val	Ala
1				5				10						15	

Val	Leu	Glu	Gly	Thr	Glu	Ser	Lys	Ile	Ile	Ala	Asn	Pro	Glu
			20				25					30	

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (A) DESCRIPTION: cBP70

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Xaa	Xaa	Xaa	Glu	Val	Ala	Lys	Xaa	Ser	Gln	Asp	Thr	Thr	Thr	Ala	Ser
1				5				10						15	

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: cBP60

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Xaa	Asn	Glu	Arg	Val	Lys	Ile	Val	Ala	Thr	Leu	Gly	Pro	Ala	Val	Glu
1				5				10				15			
Gly Arg Gly															

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: cBP50 pep

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Xaa	Ile	Ile	Xaa	Xaa	Val	Tyr	Ala	Arg	Glu	Val	Leu	Asp	Ser	Arg	Gly
1				5					10					15	
Asn Pro															

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP112-Int1

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Glu	Asp	Arg	Arg	Asn	Tyr	Pro	Thr	Asn	Thr	Tyr	Lys
1				5						10	

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(A) DESCRIPTION: cBP112-Int2

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Xaa	Asp	Asp	Gln	Gln	Ala	Glu	Glu	Asp	Tyr	Ala
1				5						10

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Trp	Gln	Pro	Pro	Arg	Ala	Arg	Ile
1				5			

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: 50 KDa protein

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *S. pneumoniae*

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ile	Thr	Asp	Val	Tyr	Ala	Arg	Glu	Val	Leu	Asp	Ser	Arg	Gly	Asn	Pro
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (A) DESCRIPTION: 50 KDa protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *B. subtilis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Ile	Val	Asp	Val	Arg	Ala	Arg	Glu	Val	Leu	Asp	Ser	Arg	Gly	Asn	Pro
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TCCCGCGGCA ACCCAACACT TGAAGTAGAA GTTTACACTG AATCAGGTGC TTTCGGACGT

GGTATGGTTC CATCAGGAGC TTCTACTGGT GAACACGAAG CAGTTGAACT TCGCGACGGT 120
GACAAATCTC GTTACGGTGG TCTTGGTACA CAAAAAGCTG TTGACAACGT AAACAACATC 180
ATTGCTGAGG CCATCATTGG CTACGATGTA 210

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: B. subtilis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TCCCGCGGCA ACCCAACAGT TGAAGTTGAA GTATATACAG AAACAGGAGC TTTCGGCCGC 60
GCATTAGTGC CAAGCGGAGC TTCTACAGGT CAATACGAAG CGGTTGAGCT TCGTGACGGC 120
GACAAAGACC GTTACCTTGG AAAAGGCGTG TTAACAGCTG TTAACAACGT AAACGAAATC 180
ATTGCTCCAG AGCTTCTTGG CTTTGATGTA 210

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Ser	Arg	Gly	Asn	Pro	Thr	Leu	Glu	Val	Glu	Val	Tyr	Thr	Glu	Ser	Gly
1				5					10					15	
Ala	Phe	Gly	Arg	Gly	Met	Val	Pro	Ser	Gly	Ala	Ser	Thr	Gly	Glu	His
				20					25					30	
Glu	Ala	Val	Glu	Leu	Arg	Asp	Gly	Asp	Lys	Ser	Arg	Tyr	Gly	Gly	Leu
				35					40					45	
Gly	Thr	Gln	Lys	Ala	Val	Asp	Asn	Val	Asn	Asn	Ile	Ile	Ala	Glu	Ala
				50					55					60	
Ile	Ile	Gly	Tyr	Asp	Val										
65															70

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *B. subtilis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Ser	Arg	Gly	Asn	Pro	Thr	Val	Glu	Val	Glu	Val	Tyr	Thr	Glu	Thr	Gly
1				5					10					15	

Ala	Phe	Gly	Arg	Ala	Leu	Val	Pro	Ser	Gly	Ala	Ser	Thr	Gly	Gln	Tyr
				20					25					30	

Glu	Ala	Val	Glu	Leu	Arg	Asp	Gly	Asp	Lys	Asp	Arg	Tyr	Leu	Gly	Lys
				35					40					45	

Gly	Val	Leu	Thr	Ala	Val	Asn	Asn	Val	Asn	Glu	Ile	Ile	Ala	Pro	Glu
				50				55						60	

Leu	Leu	Gly	Phe	Asp	Val
65					70

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1511 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: B. subtilis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TCCCGCGGCA ACCCAACACT TGAAGTAGAA GTTTACACTG AATCAGGTGC TTTCGGACGT	60
GGTATGGTTC CATCAGGAGC TTCTACTGGT GAACACGAAG CAGTTGAACT TCGCGACGGT	120
GACAAATCTC GTTACGGTGG TCTTGGTACA CAAAAGCTG TTGACAACGT AAACAACATC	180
ATTGCTGATT CTATCATTGG CTACGATGTA CGTGATCAAC AAGCTATTGA CCGTGCTATG	240
ATCGCACTTG ACGGTACTCC TAACAAAGGT AAATTGGGTG CGAATGCAAT CCTCGGTGTG	300
TCTATCGCTG TAGCTCGTGC TGCTGCTGAC TACCTTGAAA TCCCACTTTA CAGCTACCTT	360
GGTGGATTCA AACTAAAGT TCTTCCAAC CCAATGATGA ACATCATCAA CGGTGGTTCT	420
CACTCTGACG CTCCAATCGC TTTCCAAGAG TTCATGATCT TGCCAGTTGG TGCGCCAACA	480
TTTAAAGAAA CCCTTCGTTA CGGTGCTGAA ATCTTCCACG CTCTTAAGAA AATCCTTAAA	540
TCACGTGGTT TGGAAACTGC CGTAGGTGAC GAAGGTGGAT TCGCTCCTCG TTTCGAAGGA	600
ACTGAAGATG GTGTTGAAAC TATCCTTGCT GCGATTGAAG CTGCTGGATA TGTACCAGGT	660

AAAGACGTAT TTCTCGGATT TGA	TGCTGTGCT TCATCAGAAT TCTACGATAA AGAACGTAAA	720
GTTTACGACT AACTAAATT TGAAGGCGAA	GGTGCTGCTG TTCGTACATC TGCAGAACAA	780
ATCGACTACC TTGAAGAATT GGTAAACAAA	TACCCAATCA TCACTATTGA AGATGGTATG	840
GATGAAAACG ACTGGGATGG TTGGAAAGCT	CTTACTGAAC GTCTTGGTAA GAAAGTACAA	900
CTTGTTGGTG ACGACTTCTT CGTAACAAAC	ACTGACTACC TTGCACGTGG TATCCAAAAA	960
GGTGCTGCTA ACTCAATCCT TATCAAAGTT	AACCAAATCG GTACTCTTAC TGAAACTTTT	1020
GAAGCTATCG AAATGGCTAA AGAAGCTGGT	TACACTGCTG TTGTATCACA CCGTTCAGGT	1080
GAAACTGAAG ATTCAACAAT CGCTGATATC	GCGATTGCAA CTAACGCAGG ACAAATCAAG	1140
ACTGGTTCAC TTTCACGTAC AGACCGTATC	GCTAAATACA ACCAATTGCT TCGTATCGAA	1200
GACCAACTTG GTGAAGTAGC AGAATATCGT	GGATTGAAAT CATTCTACGA CCTTAAAAAA	1260
TAAATAGTA CAGTGAAC TA TTTTATCCCC	GAACCATGAA ATTCAAATTC CGGCCCTTGA	1320
TTTAACAAC TTCTAGCCCC CCGATTATC	CGAGGGGATA TTTTGT TTTT AAAGGGCAAA	1380
AAGGACAAAC TTTTAAAT CTTTTCATAC	ATTACCATAC TTGATGCTCC CCTCCTTCTT	1440
GTACTGGTTT AAGAATATAC TCCATCGATG	ACACCGTCGG ATGTTTAGGA TTATTCGACA	1500
ACAAGTTGAA T		1511

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 420 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:

(ii) MOLECULE TYPE: protein

(iv) ANTI-SENSE: NO

(A) ORGANISM: B. subtilis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ser Arg Gly Asn Pro Thr Leu Glu Val Glu Val Tyr Thr Glu Ser Gly
1 5 10 15

Ala Phe Gly Arg Gly Met Val Pro Ser Gly Ala Ser Thr Gly Glu His
20 25 30

Glu Ala Val Glu Leu Arg Asp Gly Asp Lys Ser Arg Tyr Gly Gly Leu
35 40 45

Gly Thr Gln Lys Ala Val Asp Asn Val Asn Asn Ile Ile Ala Asp Ser
50 55 60

Ile Ile Gly Tyr Asp Val Arg Asp Gln Gln Ala Ile Asp Arg Ala Met
65 70 75 80

Ile Ala Leu Asp Gly Thr Pro Asn Lys Gly Lys Leu Gly Ala Asn Ala
85 90 95

Ile Leu Gly Val Ser Ile Ala Val Ala Arg Ala Ala Ala Asp Tyr Leu
100 105 110

Glu Ile Pro Leu Tyr Ser Tyr Leu Gly Gly Phe Asn Thr Lys Val Leu
115 120 125

Gly Ala Ala Asn Ser Ile Leu Ile Lys Val Asn Gln Ile Gly Thr Leu

325

330

335

Thr Glu Thr Phe Glu Ala Ile Glu Met Ala Lys Glu Ala Gly Tyr Thr
 340 345 350

Ala Val Val Ser His Arg Ser Gly Glu Thr Glu Asp Ser Thr Ile Ala
 355 360 365

Asp Ile Ala Ile Ala Thr Asn Ala Gly Gln Ile Lys Thr Gly Ser Leu
 370 375 380

Ser Arg Thr Asp Arg Ile Ala Lys Tyr Asn Gln Leu Leu Arg Ile Glu
 385 390 395 400

Asp Gln Leu Gly Glu Val Ala Glu Tyr Arg Gly Leu Lys Ser Phe Tyr
 405 410 415

Asp Leu Lys Lys
 420

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 429 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

AAAAAAGGTA GCAGAAGCTG AGAAGAAGGT TGAAGAAGCT GAGAAAAAAG CCAAGGATCA	60
AAAAGAAGAA GATCGCCGTA ACTACCCAAC CAATACTTAC AAAACGCTTG ACCTTGAAAT	120
TGCTGAGTCC GATGTGAAAG TTAAAGAAGC GGAGCTTGAA CTAGTAAAAG AGGAAGCTAA	180
GGAACCTCGA GACGAGGAAA AAATTAAGCA AGCAAAAGCG AAAGTTGAGA GTAAAAAAGC	240
TGAGGCTACA AGGTTAGAAA ACATCAAGAC AGATCGTAAA AAAGCAGAAG AAGAAGCTAA	300
ACGAAAAGCA GCAGAAGAAG ATAAAGTTAA AGAAAAACCA GCTGAACAAC CACAACCAGC	360
GCCGGTTACT CAACCAGAAA AACCAGCTCC AAAACCAGAG AAGCCAGCTG AACCAACCAA	420
AGCAGAAAA	429

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 142 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *S. pneumoniae*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu Glu Ala Glu Lys Lys
 1 5 10 15
 Ala Lys Asp Gln Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Asn Thr
 20 25 30
 Tyr Lys Thr Leu Asp Leu Glu Ile Ala Glu Ser Asp Val Lys Val Lys
 35 40 45
 Glu Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys Glu Pro Arg Asp
 50 55 60
 Glu Glu Lys Ile Lys Gln Ala Lys Ala Lys Val Glu Ser Lys Lys Ala
 65 70 75 80
 Glu Ala Thr Arg Leu Glu Asn Ile Lys Thr Asp Arg Lys Lys Ala Glu
 85 90 95
 Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp Lys Val Lys Glu Lys
 100 105 110
 Pro Ala Glu Gln Pro Gln Pro Ala Pro Val Thr Gln Pro Glu Lys Pro
 115 120 125
 Ala Pro Lys Pro Glu Lys Pro Ala Glu Gln Pro Lys Ala Glu
 130 135 140

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide primer"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GCTCTTNCTC GATGTCTCNG TNGCCAT

27

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide primer"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AGCATAGTCT TCTTCGACTT GTTGATCATC

30

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2462 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCAAGCAGCC ACTTCTTCTA ATATGGCAAA GACAGAACAT AGGAAAGCTG CTAACAAGC	60
GTCGATGAAT ATATAGAAAA AATGATGAGG GAGATTCAAC TAGATAGAAG AAAACATACC	120
CAAAATGTCG CCTTAAACAT AAAGTTGAGC GCAATTAAAA CGAAGTATTT GCGTGAATTA	180
AATGTTTTAG AAGAGAAGTC GAAAGATGAG TTGCCGTCAG AAATAAAAGC AAAGTTAGAC	240
GCAGCTTTTG AGAAGTTTAA AAAAGATACA TTGAAACCAG GAGAAAAGGT AGCAGAAGCT	300
AAGAAGAAGG TTGAAGAAGC TAAGAAAAAA GCCGAGGATC AAAAGAAGA AGATCGTCGT	360
AACTACCCAA CCAATACTTA CAAAACGCTT GAACTTGAAA TTGCTGAGTT CGATGTGAAA	420
GTTAAAGAAG CGGAGCTTGA ACTAGTAAAA GAGGAAGCTA AAGAATCTCG AAACGAGGGC	480
ACAATTAAGC AAGCAAAAGA GAAAGTTGAG AGTAAAAAAG CTGAGGCTAC AAGGTTAGAA	540
AACATCAAGA CAGATCGTAA AAAAGCAGAA GAAGAAGCTA AACGAAAAGC AGATGCTAAG	600
TTGAAGGAAG CTAATGTAGC GACTTCAGAT CAAGGTAAAC CAAAGGGGCG GGCAAAACGA	660

GGAGTTCCTG	GAGAGCTAGC	AACACCTGAT	AAAAAAGAAA	ATGATGCGAA	GTCTTCAGAT	720
TCTAGCGTAG	GTGAAGAAAC	TCTTCCAAGC	TCATCCCTGA	AATCAGGAAA	AAAGGTAGCA	780
GAAGCTGAGA	AGAAGGTTGA	AGAAGCTGAG	AAAAAAGCCA	AGGATCAAAA	AGAAGAAGAT	840
CGCCGTAACT	ACCCAACCAA	TACTTACAAA	ACGCTTGACC	TTGAAATTGC	TGAGTCCGAT	900
GTGAAAGTTA	AAGAAGCGGA	GCTTGAACTA	GTAAAAGAGG	AAGCTAAGGA	ACCTCGAGAC	960
GAGGAAAAAA	TTAAGCAAGC	AAAAGCGAAA	GTTGAGAGTA	AAAAAAGCTGA	GGCTACAAGG	1020
TTAGAAAACA	TCAAGACAGA	TCGTAAAAAA	GCAGAAGAAG	AAGCTAAACG	AAAAGCAGCA	1080
GAAGAAGATA	AAGTTAAAGA	AAAACCAGCT	GAACAACCAC	AACCAGCGCC	GGCTACTCAA	1140
CCAGAAAAAC	CAGCTCCAAA	ACCAGAGAAG	CCAGCTGAAC	AACCAAAAGC	AGAAAAAACA	1200
GATGATCAAC	AAGCTGAAGA	AGACTATGCT	CGTAGATCAG	AAGAAGAATA	TAATCGCTTG	1260
ACTCAACAGC	AACCGCCAAA	AACTGAAAAA	CCAGCACAAAC	CATCTACTCC	AAAAACAGGC	1320
TGGAAACAAG	AAAACGGTAT	GTGGTACTTC	TACAATACTG	ATGGTTCAAT	GGCAACAGGA	1380
TGGCTCCAAA	ACAACGGTTC	ATGGTACTAT	CTAAACGCTA	ATGGTGCTAT	GGCGACAGGA	1440
TGGCTCCAAA	ACAATGGTTC	ATGGTACTAT	CTAAACGCTA	ATGGTTCAAT	GGCAACAGGA	1500
TGGCTCCAAA	ACAATGGTTC	ATGGTACTAC	CTAAACGCTA	ATGGTGCTAT	GGCGACAGGA	1560
TGGCTCCAAT	ACAATGGTTC	ATGGTACTAC	CTAAACAGCA	ATGGCGCTAT	GGCGACAGGA	1620
TGGCTCCAAT	ACAATGGCTC	ATGGTACTAC	CTCAACGCTA	ATGGTGATAT	GGCGACAGGA	1680
TGGCTCCAAA	ACAACGGTTC	ATGGTACTAC	CTCAACGCTA	ATGGTGATAT	GGCGACAGGA	1740
TGGCTCCAAT	ACAACGGTTC	ATGGTATTAC	CTCAACGCTA	ATGGTGATAT	GGCGACAGGT	1800

TGGGTGAAAG ATGGAGATAC CTGGTACTAT CTTGAAGCAT CAGGTGCTAT GAAAGCAAGC	1860
CAATGGTTCA AAGTATCAGA TAAATGGTAC TATGTCAATG GCTCAGGTGC CCTTGCAGTC	1920
AACACAACCTG TAGATGGCTA TGGAGTCAAT GCCAATGGTG AATGGGTAAA CTAAACCTAA	1980
TATAACTAGT TAATACTGAC TTCCTGTAAG AACTTTTTTAA AGTATTCCCT ACAAATACCA	2040
TATCCTTTCA GTAGATAATA TACCCTTGTA GGAAGTTTAG ATTAAAAAAT AACTCTGTAA	2100
TCTCTAGCCG GATTTATAGC GCTAGAGACT ACGGAGTTTT TTTGATGAGG AAAGAATGGC	2160
GGCATTCAAG AGACTCTTTA AGAGAGTTAC GGGTTTTTAAA CTATTAAGCC TTCTCCAATT	2220
GCAAGAGGCT TCAATCTCTG CTAGGGTGCT AGCTTGCGAA ATGGCTCCAC GGAGTTTGGC	2280
AGCGCCAGAT GTTCCACGGA GATAGTGAGG AGCGAGGCCG CGGAATTCAC GAACTGCGAC	2340
GTTTTCTCCT TTGAGGTAA TCAATCGTTT CAAGTGTTTC TAGGCGATCT TCATCTTGTC	2400
TTCAAAGGTC AAATCAGGTA GGATTTCTCC TGTTTCAAAG TTTATGGTGG CCCTGGTTGA	2460
AG	2462

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 631 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Met	Met	Arg	Glu	Ile	Gln	Leu	Asp	Arg	Arg	Lys	His	Thr	Gln	Asn	Val
1				5					10					15	
Ala	Leu	Asn	Ile	Lys	Leu	Ser	Ala	Ile	Lys	Thr	Lys	Tyr	Leu	Arg	Glu
			20					25					30		
Leu	Asn	Val	Leu	Glu	Glu	Lys	Ser	Lys	Asp	Glu	Leu	Pro	Ser	Glu	Ile
		35					40					45			
Lys	Ala	Lys	Leu	Asp	Ala	Ala	Phe	Glu	Lys	Phe	Lys	Lys	Asp	Thr	Leu
	50					55					60				
Lys	Pro	Gly	Glu	Lys	Val	Ala	Glu	Ala	Lys	Lys	Lys	Val	Glu	Glu	Ala
65					70					75				80	
Lys	Lys	Lys	Ala	Glu	Asp	Gln	Lys	Glu	Glu	Asp	Arg	Arg	Asn	Tyr	Pro
			85						90					95	
Thr	Asn	Thr	Tyr	Lys	Thr	Leu	Glu	Leu	Glu	Ile	Ala	Glu	Phe	Asp	Val
			100						105					110	
Lys	Val	Lys	Glu	Ala	Glu	Leu	Glu	Leu	Val	Lys	Glu	Glu	Ala	Lys	Glu
		115						120					125		
Ser	Arg	Asn	Glu	Gly	Thr	Ile	Lys	Gln	Ala	Lys	Glu	Lys	Val	Glu	Ser
		130					135					140			
Lys	Lys	Ala	Glu	Ala	Thr	Arg	Leu	Glu	Asn	Ile	Lys	Thr	Asp	Arg	Lys
145					150					155				160	
Lys	Ala	Glu	Glu	Glu	Ala	Lys	Arg	Lys	Ala	Asp	Ala	Lys	Leu	Lys	Glu
				165					170					175	

Ala Asn Val Ala Thr Ser Asp Gln Gly Lys Pro Lys Gly Arg Ala Lys
 180 185 190

Arg Gly Val Pro Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu Asn Asp
 195 200 205

Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Pro Ser Ser
 210 215 220

Ser Leu Lys Ser Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu
 225 230 235 240

Glu Ala Glu Lys Lys Ala Lys Asp Gln Lys Glu Glu Asp Arg Arg Asn
 245 250 255

Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Asp Leu Glu Ile Ala Glu Ser
 260 265 270

Asp Val Lys Val Lys Glu Ala Glu Leu Glu Leu Val Lys Glu Glu Ala
 275 280 285

Lys Glu Pro Arg Asp Glu Glu Lys Ile Lys Gln Ala Lys Ala Lys Val
 290 295 300

Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Asn Ile Lys Thr Asp
 305 310 315 320

Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp
 325 330 335

Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Thr
 340 345 350

Gln Pro Glu Lys Pro Ala Pro Lys Pro Glu Lys Pro Ala Glu Gln Pro
 355 360 365

Lys Ala Glu Lys Thr Asp Asp Gln Gln Ala Glu Glu Asp Tyr Ala Arg

370	375	380
Arg Ser Glu Glu Glu Tyr Asn Arg Leu Thr Gln Gln Gln Pro Pro Lys		
385	390	395 400
Thr Glu Lys Pro Ala Gln Pro Ser Thr Pro Lys Thr Gly Trp Lys Gln		
405	410	415
Glu Asn Gly Met Trp Tyr Phe Tyr Asn Thr Asp Gly Ser Met Ala Thr		
420	425	430
Gly Trp Leu Gln Asn Asn Gly Ser Trp Tyr Tyr Leu Asn Ala Asn Gly		
435	440	445
Ala Met Ala Thr Gly Trp Leu Gln Asn Asn Gly Ser Trp Tyr Tyr Leu		
450	455	460
Asn Ala Asn Gly Ser Met Ala Thr Gly Trp Leu Gln Asn Asn Gly Ser		
465	470	475 480
Trp Tyr Tyr Leu Asn Ala Asn Gly Ala Met Ala Thr Gly Trp Leu Gln		
485	490	495
Tyr Asn Gly Ser Trp Tyr Tyr Leu Asn Ser Asn Gly Ala Met Ala Thr		
500	505	510
Gly Trp Leu Gln Tyr Asn Gly Ser Trp Tyr Tyr Leu Asn Ala Asn Gly		
515	520	525
Asp Met Ala Thr Gly Trp Leu Gln Asn Asn Gly Ser Trp Tyr Tyr Leu		
530	535	540
Asn Ala Asn Gly Asp Met Ala Thr Gly Trp Leu Gln Tyr Asn Gly Ser		
545	550	555 560
Trp Tyr Tyr Leu Asn Ala Asn Gly Asp Met Ala Thr Gly Trp Val Lys		
565	570	575

Asn Gly Glu Trp Val Asn Glx
625 630